

Anti-EXOSC2 antibody (1-293) (STJ112479)
STJ112479

GENERAL INFORMATION


Product Type	Primary antibodies
Short Description	
Applications	WB/IF/ICC/ELISA
Host/Source	Rabbit
Reactivity	Human/Mouse/Rat

PRODUCT PROPERTIES


Clonality	Polyclonal
Clone ID	
Concentration	Lot specific
Conjugation	Unconjugated
Purification	Affinity purification
Dilution Range	WB: 1:500-1:2000 IF/CC: 1:50-1:200
	ELISA: Recommended starting concentration is 1 μ g/mL. Please optimize the concentration based on your specific assay requirements.
Formulation	PBS with 0.01% Thimerosal, 50% Glycerol, pH 7.3.
Isotype	IgG
Storage Instruction	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

TARGET INFORMATION


Gene ID	23404
Gene Symbol	EXOSC2
Uniprot ID	EXOS2_HUMAN
Immunogen Region	1-293
Specificity	Recombinant fusion protein containing a sequence corresponding to amino acids 1-293 of human EXOSC2 (NP_055100.2).
Immunogen Sequence	MAMEMRLPVARKPLSERLGR DTKKHLVVPDITTTDTGFM RGHGTYMGEEKLIASVAGSV ERVNLKICVKALKTRYIGEV GDIVVGRITEVQQKRWKVET NSRLDSVLLSSMNLPGGEL RRRSAEDELAMRGFLQEGDL ISAEVQAVFSDGAVSLHTRS LKYGKLGQGVLVQVSPSLVK RQKTHFDLPCGASVILGNN GFIWIYPTPEHKEEEAGGFI ANLEPVSLADREVISRLRN




Immunofluorescence analysis of H9C2 cells using EXOSC2 Rabbit polyclonal antibody (STJ112479) at dilution of 1:100. Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of L929 cells using EXOSC2 Rabbit polyclonal antibody (STJ112479) at dilution of 1:100. Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of U2OS cells using EXOSC2 Rabbit polyclonal antibody (STJ112479) at dilution of 1:100. Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) at 1:500 dilution. Blue: DAPI for nuclear staining.



Western blot analysis of various lysates using EXOSC2 Rabbit polyclonal antibody (STJ112479) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJ5000856) at 1:10000 dilution. Lysates/proteins: 25 μ g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 5s.

This product is suitable for in-vitro studies under the RESEARCH USE ONLY [RUO] licence. This product must not be used as for diagnostic or other medical purposes.
St John's Laboratory Ltd, Knowledge Dock Business Centre, University Way, London, E16 2RD | Tel: 0208 223 3081