

**Anti-SERPINI2 antibody (56-405) (STJ113329)**  
STJ113329

**GENERAL INFORMATION**


<b>Product Type</b>	Primary antibodies
<b>Short Description</b>	
<b>Applications</b>	WB/IF/ICC/ELISA
<b>Host/Source</b>	Rabbit
<b>Reactivity</b>	Mouse/Rat

**PRODUCT PROPERTIES**


<b>Clonality</b>	Polyclonal
<b>Clone ID</b>	
<b>Concentration</b>	Lot specific
<b>Conjugation</b>	Unconjugated
<b>Purification</b>	Affinity purification
<b>Dilution</b>	WB:1:500-1:2000
<b>Range</b>	IF/CC:1:50-1:100 ELISA:Recommended starting concentration is 1 $\mu$ g/mL. Please optimize the concentration based on your specific assay requirements.
<b>Formulation</b>	PBS with 0.01% Thimerosal, 50% Glycerol, pH 7.3.
<b>Isotype</b>	IgG
<b>Storage Instruction</b>	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

**TARGET INFORMATION**


<b>Gene ID</b>	5276
<b>Gene Symbol</b>	SERPINI2
<b>Uniprot ID</b>	SPI2_HUMAN
<b>Immunogen</b>	
<b>Immunogen Region</b>	56-405
<b>Specificity</b>	Recombinant fusion protein containing a sequence corresponding to amino acids 56-405 of human SERPINI2 (NP_006208.1).
<b>Immunogen Sequence</b>	LEMVQLGAKGKAQQIRQTL KQQETSAGEEFFVLKSFSA ISEKKQEFTNLNANALYLQE GFTVKEQYLHGNKEFFQSAI KLVDFAQDAKACAEMISTWVE RKTGDKIKDMFSGEEFGPLT RLVLVNAIYFKGDWKQKFRK EDTQLINFTKKNGSTVKIPM MKALLRTKYGYFSESSLNYQ VLELSYKGDSEFLIILPAE GMDIEEVEKLITAAQILKWL SEMQEEVEISLPRFKVEQ



Immunofluorescence analysis of paraffin-embedded mouse pancreas using SERPINI2 Rabbit polyclonal antibody (STJ113329) at dilution of 1:100 (40x lens). Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of paraffin-embedded rat pancreas using SERPINI2 Rabbit polyclonal antibody (STJ113329) at dilution of 1:100 (40x lens). Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) at 1:500 dilution. Blue: DAPI for nuclear staining.



Western blot analysis of various lysates using SERPINI2 Rabbit polyclonal antibody (STJ113329) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJ500856) at 1:10000 dilution. Lysates/proteins: 25  $\mu$ g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 10s.

This product is suitable for in-vitro studies under the RESEARCH USE ONLY [RUO] licence. This product must not be used as for diagnostic or other medical purposes.  
St John's Laboratory Ltd, Knowledge Dock Business Centre, University Way, London, E16 2RD | Tel: 0208 223 3081