

Anti-CLU antibody (314-449) (STJ114779)

STJ114779

GENERAL INFORMATION

Product Type Primary antibodies

Short

Description
Applications WR/III

Applications WB/IHC-P/ELISA

Host/Source Rabbit
Reactivity Human/Mouse/Rat

PRODUCT PROPERTIES

Clonality Polyclonal

Clone ID

Concentration Lot specific

Conjugation
Purification
Dilution
Range
Unconjugated
Affinity purification
WB:1:500-1:1000
HC-P:1:50-1:200

ELISA:Recommended starting concentration is 1 Mu g/mL. Please optimize the concentration based on your specific assay

requirements.

Formulation PBS with 0.01% Thimerosal, 50% Glycerol, pH 7.3.

Isotype IgG

Storage Store at-20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

Instruction

TARGET INFORMATION

Gene ID 1191

Gene Symbol CLU

Uniprot ID CLUS_HUMAN

Immunogen

Immunogen 314-449


Region Specificity

Recombinant fusion protein containing a sequence corresponding to amino acids 314-449 of human Clusterin alpha chain


(NP_001822.3).

 $\textbf{Immunogen} \quad \textbf{STNNPSQAKLRRELDESLQV} \text{ AERLTRKYNELLKSYQWKML NTSSLLEQLNEQFNWVSRLA NLTQGEDQYYLRVTTVASHT} \\$


Sequence SDSDVPSGVTEVVVKLFDSD PITVTVPVEVSRKNPKFMET VAEKALQEYRKKHREE




Immunohistochemistry analysis of Clusterin alpha chai in paraffin-embedded human esophageal usin Clusterin alpha chain Rabbit polyclonal antibod (STJ114779) at dilution of 1:100 (40x lens). Perform hig pressure antigen retireval with 10 mM citrate buffer pl 6. 0 before commencing with immunohistochemistr



Immunohistochemistry analysis of Clusterin alpha chair in paraffin-embedded human colon carcinoma using Clusterin alpha chair Rabbit polyclonal antibody (STJ114779) at dilution of 1:100 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pl-6. 0 before commencing with immunohistochemistry.



Immunohistochemistry analysis of Clusterin alpha chai in paraffin-embedded human lung using Clusterin alph chain Rabbit polyclonal antibody (STJ114779) dilution of 1:100 (40x lens). Perform high pressu antigen retrieval with 10 mM citrate buffer pH 6. before commencing with immunohistochemist



Immunohistochemistry analysis of Clusterin alpha chain in paraffin-embedded rat spleen using Clusterin alpha chain Rabbit polyclonal antibody (STJ114779) at dilution of 1:100 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6. 0 before commencing with immunohistochemistry stalining protocol.