

Anti-DNAJB11 antibody (89-358) (STJ114781)
STJ114781

GENERAL INFORMATION


Product Type	Primary antibodies
Short Description	
Applications	WB/IHC-P/ELISA
Host/Source	Rabbit
Reactivity	Human/Mouse/Rat

PRODUCT PROPERTIES


Clonality	Polyclonal
Clone ID	
Concentration	Lot specific
Conjugation	Unconjugated
Purification	Affinity purification
Dilution Range	WB:1:500-1:2000 IHC-P:1:50-1:200 ELISA:Recommended starting concentration is 1 Mu g/mL. Please optimize the concentration based on your specific assay requirements.
Formulation	PBS with 0.01% Thimerosal, 50% Glycerol, pH 7.3.
Isotype	IgG
Storage Instruction	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

TARGET INFORMATION


Gene ID	51726
Gene Symbol	DNAJB11
Uniprot ID	DJB11_HUMAN
Immunogen	
Immunogen Region	89-358
Specificity	Recombinant fusion protein containing a sequence corresponding to amino acids 89-358 of human DNAJB11 (NP_057390.1).
Immunogen Sequence	YGEGLKDGHQSSHGDIFFSH FFGDFGFMFGGTPRQQDRNI PRGSDIIVLEVTLEEVYAG NFVEVRNKPVARQAPGKRK CNCRQEMRTTQLGPGRFQMT QEVWCDECPNVKLVNEERTL EVEIEPGVRDGMIEYPIGEG EPHVDGEPGDLRFRIKVVKH PIFERRGGDLYTNVTISLVE SLVGFEMDITHLDGHKHHIS RDKITRPGAKLWKKGEGLPN FDNNNIKGSLITFDVDFP




Immunohistochemistry analysis of paraffin-embedded mouse spinal cord using DNAJB11 antibody (STJ114781) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with immunohistochemistry staining protocol.



Immunohistochemistry analysis of paraffin-embedded rat brain using DNAJB11 antibody (STJ114781) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with immunohistochemistry staining protocol.



Immunohistochemistry analysis of paraffin-embedded rat testis using DNAJB11 antibody (STJ114781) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with immunohistochemistry staining protocol.



Western blot analysis of extracts of various cell lines using DNAJB11 antibody (STJ114781) at 1:3000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJ5000856) at 1:10000 dilution. Lysates/proteins: 25 Mu g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 90s.

This product is suitable for in-vitro studies under the RESEARCH USE ONLY [RUO] licence. This product must not be used as for diagnostic or other medical purposes.
St John's Laboratory Ltd, Knowledge Dock Business Centre, University Way, London, E16 2RD | Tel: 0208 223 3081