

Anti-TBL1XR1 antibody (120-170) (STJ115399)

STJ115399

GENERAL INFORMATION

Product Type Primary antibodies

Short Description

Applications WB/IF/ICC/ELISA
Host/Source Rabbit
Reactivity Human/Mouse

PRODUCT PROPERTIES

Clonality Polyclonal

Clone ID

Concentration
Conjugation
Purification
Dilution Range
WB:1:500-1::1000
WB:1:500-1::1000

IF/ICC:1:50-1:200

ELISA:Recommended starting concentration is 1 Mu g/mL. Please optimize the concentration based on your specific assay

requirements.

Formulation PBS with 0.09% Sodium Azide, 50% Glycerol, pH 7.3.

Isotype IgG

Storage Instruction Store at-20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

TARGET INFORMATION

Gene ID 79718
Gene Symbol TBL1XR1
Uniprot ID TBL1R_HUMAN


Immunogen

Immunogen Region 120-170


Specificity Recombinant fusion protein containing a sequence corresponding to amino acids 120-170 of human TBL1XR1 (NP_078941.2).

Immunogen QQGSAKNGENTANGEENGAH TIANNHTDMMEVDGDVEIPP NKAVVLRGHES


Sequence



Immunofluorescence analysis of A549 cells using TBL1XR1 antibody (STJ115399).



Western blot analysis of extracts of Mouse brain, usin TBL1XR1 antibody (STJ115399) at 1:1000 dilutior Secondary antibody: HRP Goat Anti-Rabbit IgG (H+I (STJS000856) at 1:10000 dilution. Lysates/proteins: 2: Mu g per lane. Blocking buffer: 3% norfat dry milk it TBST. Detection: ECL Enhanced Kit. Exposure time 180s.



Western blot analysis of extracts of K-562 cells, using TBL1XR1 antibody (STJ115399) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJS000566) at 1:10000 dilution. Lysates/proteins-L) 5 Mu g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit Exposure time: 1805.