

Anti-ACY1 antibody (1-408) (STJ115419)

STJ115419

GENERAL INFORMATION


Product Type	Primary antibodies
Short Description	
Applications	WB/IHC-P/IF/ICC/ELISA
Host/Source	Rabbit
Reactivity	Human/Mouse/Rat

PRODUCT PROPERTIES


Clonality	Polyclonal
Clone ID	
Concentration	Lot specific
Conjugation	Unconjugated
Purification	Affinity purification
Dilution	WB:1:500-1:2000
Range	IHC-P:1:50-1:200 IF/ICC:1:50-1:200 ELISA:Recommended starting concentration is 1 μ g/mL. Please optimize the concentration based on your specific assay requirements.
Formulation	PBS with 0.02% Sodium Azide, 50% Glycerol, pH 7.3.
Isotype	IgG
Storage	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.
Instruction	

TARGET INFORMATION


Gene ID	95
Gene Symbol	ACY1
Uniprot ID	ACY1_HUMAN
Immunogen	
Immunogen Region	1-408
Specificity	Recombinant fusion protein containing a sequence corresponding to amino acids 1-408 of human ACY1 (NP_001185824.1).
Immunogen Sequence	MTSKGPEEEEHPSVTLFRQYL RIRTVQPKPDYGAAVAFFEE TARQLGLGCQKVEVAPGYV TVLTWPGTNPTLSSILLNSH TDWVPVFKEHWSHDPFEAFK DSEGYIYARGAQDMKCVSIQ YLEAVRRLLKVEGHRFPRTIH MTFVPDEEVGGHQQGMELFVQ RPEFHALRAGFALDEGIANP TDAFTVFYSERSPWVVRVTS TGRPGHASRFMEDTAAEKLH KVVNSILAFREKEWQRLQS




Immunofluorescence analysis of MCF7 cells using ACY1 antibody (STJ115419). Blue: DAPI for nuclear staining.



Immunohistochemistry analysis of paraffin-embedded human esophageal cancer using ACY1 antibody (STJ115419) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with immunohistochemistry staining protocol.



Immunohistochemistry analysis of paraffin-embedded rat kidney using ACY1 antibody (STJ115419) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with immunohistochemistry staining protocol.



Western blot analysis of extracts of various cell lines, using ACY1 antibody (STJ115419) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJ/S000856) at 1:10000 dilution. Lysates/proteins: 25 μ g per lane. Blocking buffer: 3% nonfat dry milk in TBST.

This product is suitable for in-vitro studies under the RESEARCH USE ONLY [RUO] licence. This product must not be used as for diagnostic or other medical purposes.
St John's Laboratory Ltd, Knowledge Dock Business Centre, University Way, London, E16 2RD | Tel: 0208 223 3081