

## Anti-IAH1 antibody (1-248) (STJ115664)

STJ115664

### GENERAL INFORMATION


|                          |                    |
|--------------------------|--------------------|
| <b>Product Type</b>      | Primary antibodies |
| <b>Short Description</b> |                    |
| <b>Applications</b>      | WB/ELISA           |
| <b>Host/Source</b>       | Rabbit             |
| <b>Reactivity</b>        | Human/Mouse/Rat    |

### PRODUCT PROPERTIES

|                      |   |
|----------------------|---|
| <b>Clonality</b>     | Polyclonal  |
| <b>Clone ID</b>      |   |
| <b>Concentration</b> | Lot specific  |
| <b>Conjugation</b>   | Unconjugated  |
| <b>Purification</b>  | Affinity purification   |
| <b>Dilution</b>      | WB:1:500-1:1000   |
| <b>Range</b>         | ELISA:Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements. |
| <b>Formulation</b>   | PBS with 0.01% Thimerosal, 50% Glycerol, pH 7.3.  |
| <b>Isotype</b>       | IgG   |
| <b>Storage</b>       | Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.                                    |
| <b>Instruction</b>   |   |

### TARGET INFORMATION

|                           |  |
|---------------------------|--|
| <b>Gene ID</b>            | 285148   |
| <b>Gene Symbol</b>        | IAH1   |
| <b>Uniprot ID</b>         | IAH1_HUMAN   |
| <b>Immunogen</b>          |  |
| <b>Immunogen Region</b>   | 1-248  |
| <b>Specificity</b>        | Recombinant fusion protein containing a sequence corresponding to amino acids 1-248 of human IAH1 (NP_001034702.1).  |
| <b>Immunogen Sequence</b> | MALCEAAGCGSALLWPRLLL FGDSITQFSFQQGGWGASLA DRLVRKCDVLRGFSGYNTR WAKIILPRLIRKGNSLDIPV<br>AVTIFFGANDSALKDENPKQ HIPLEEYAANLKSVMQYLKS VDIPENRVILITPTPLCETA WEEQCIQGGCKLNRLNSVVG<br>EYANAQLQVAQDCGTDVLDL WTLMQDSQDFSSYLSDDLHL SPKGNEFLFSLWPLIEKKV SSLPLLLPYWRDVAEAKPE |



Western blot analysis of various lysates, using IAH1 Rabbit polyclonal antibody (STJ115664) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 3s.

This product is suitable for in-vitro studies under the RESEARCH USE ONLY [RUO] licence. This product must not be used as for diagnostic or other medical purposes.  
St John's Laboratory Ltd, Knowledge Dock Business Centre, University Way, London, E16 2RD | Tel: 0208 223 3081