

Anti-VCL antibody (751-850) (STJ116126)

GENERAL INFORMATION

Product Type Primary antibodies

Short Description

Applications WB/IF/ICC/ELISA Host/Source Rabbit Reactivity Human/Mouse/Rat

PRODUCT PROPERTIES

Clonality Polyclonal

Clone ID Concentration Lot specific Conjugation Unconjugated Purification Affinity purification Dilution Range WB:1:500-1:1000

IF/ICC:1:50-1:200

ELISA:Recommended starting concentration is 1 Mu g/mL. Please optimize the concentration based on your specific assay

requirements.

Formulation PBS with 0.05% Proclin300, 50% Glycerol, pH 7.3.

Isotype IaG

Storage Instruction Store at-20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

TARGET INFORMATION

Gene ID 7414

Gene Symbol VCL


Uniprot ID VINC_HUMAN

Immunogen


Immunogen 751-850

Region
Specificity
Recombinant fusion protein containing a sequence corresponding to amino acids 751-850 of human Vinculin (NP_054706.1). Immunogen AGATSIARRANRILLVAKRE VENSEDPKFREAVKAASDEL SKTISPMVMDAKAVAGNISD PGLQKSFLDSGYRILGAVAK


Sequence VREAFQPQEPDFPPPPDLE




Immunofluorescence analysis of NIH/3T3 cells using Vinculin antibody (STJ116126) at dilution of 1:150. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of U2OS cells using Vinculin antibody (STJ116126) at dilution of 1:150. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of PC-12 cells using Vinculin antibody (STJ116126) at dilution of 1:150. Blue: DAPI for nuclear staining.



Western blot analysis of extracts of various cell lines, using Vinculin antibody (STJ116126) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H-L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25 Mu g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 10s.