

Anti-MAGEA6 antibody (75-314) (STJ116318)
STJ116318

GENERAL INFORMATION


Product Type	Primary antibodies
Short Description	
Applications	WB/IF/ICC/ELISA
Host/Source	Rabbit
Reactivity	Human/Mouse

PRODUCT PROPERTIES


Clonality	Polyclonal
Clone ID	
Concentration	Lot specific
Conjugation	Unconjugated
Purification	Affinity purification
Dilution Range	WB:1:500-1:2000 IF/ICC:1:50-1:200 ELISA:Recommended starting concentration is 1 Mu g/mL. Please optimize the concentration based on your specific assay requirements.
Formulation	PBS with 0.02% Sodium Azide, 50% Glycerol, pH 7.3.
Isotype	IgG
Storage Instruction	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

TARGET INFORMATION


Gene ID	4105
Gene Symbol	MAGEA6
Uniprot ID	MAGA6_HUMAN
Immunogen Region	75-314
Specificity	Recombinant fusion protein containing a sequence corresponding to amino acids 75-314 of human MAGEA6 (NP_005354.1).
Immunogen Sequence	MNYPLWSQSYEDSSNQEEEG PSTFPDLESEFQAALSRKVA KLVHFLLLKYRAREPVTKAE MLGSVVGNGWQYFFPVIFSKA SDSLQLVFGIELMEVDPIGH VYIFATCLGLSYDGLLDNQ IMPKTGFLIILAIAGEGD CAPEEKIWEELSVLEVFEGR EDSIFGDPKKLLTQYFVQEN YLEYRQVPGSDPACYEFLWG PRALIETSYVKVLHMHVKIS GGPRISYPLLHEWALREGE




Western blot analysis of extracts of various cell lines, using MAGEA6 antibody (STJ116318) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% non-fat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 90s.



Immunofluorescence analysis of U2OS cells using MAGEA6 antibody (STJ116318) at dilution of 1:100. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of U2OS cells using MAGEA6 antibody (STJ116318) at dilution of 1:100. Blue: DAPI for nuclear staining.



Western blot analysis of extracts of various cell lines, using MAGEA6 antibody (STJ116318) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJ5000856) at 1:10000 dilution. Lysates/proteins: 25 Mu g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 90s.

This product is suitable for in-vitro studies under the RESEARCH USE ONLY [RUO] licence. This product must not be used as for diagnostic or other medical purposes.
St John's Laboratory Ltd, Knowledge Dock Business Centre, University Way, London, E16 2RD | Tel: 0208 223 3081