

## Anti-Phospho-ULK1-S757 antibody (STJ116374)

STJ116374

### GENERAL INFORMATION

<b>Product Type</b>	Primary antibodies
<b>Short Description</b>	WB/ELISA
<b>Applications</b>	Rabbit
<b>Host/Source</b>	Human
<b>Reactivity</b>	

### PRODUCT PROPERTIES

<b>Clonality</b>	Polyclonal
<b>Clone ID</b>	
<b>Concentration</b>	Lot specific
<b>Conjugation</b>	Unconjugated
<b>Purification</b>	Affinity purification
<b>Dilution Range</b>	WB:1:100-1:500 ELISA:Recommended starting concentration is 1 Mu g/mL. Please optimize the concentration based on your specific assay requirements.
<b>Formulation</b>	PBS with 0.01% Thimerosal, 50% Glycerol, pH 7.3.
<b>Isotype</b>	IgG
<b>Storage Instruction</b>	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

### TARGET INFORMATION

<b>Gene ID</b>	8408
<b>Gene Symbol</b>	ULK1
<b>Uniprot ID</b>	ULK1_HUMAN
<b>Immunogen</b>	TVGSP
<b>Immunogen Region</b>	
<b>Specificity</b>	A synthetic phosphorylated peptide around S757 of human Phospho-ULK1-S757 (NP_003556.1).
<b>Immunogen</b>	TVGSP
<b>Sequence</b>	



Western blot analysis of extracts of A-431 cells, using Phospho-ULK1-S757 antibody (STJ116374) at 1:500 dilution. A-431 cells were treated by EGF (100 ng/ml) at 37 °C for 30 minutes after serum starvation overnight. Secondary antibody: Goat Anti-Rabbit IgG (H+L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25 μg per lane. Blocking buffer: 3% non-fat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 180s.

Western blot analysis of lysates from A-431 cells, using Phospho-ULK1-S757 antibody (STJ116374) at 1:500 dilution. A-431 cells were treated by EGF (100 ng/ml) at 37 °C for 30 minutes after serum starvation overnight. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25 μg per lane. Blocking buffer: 3% non-fat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 180s.

This product is suitable for in-vitro studies under the RESEARCH USE ONLY [RUO] licence. This product must not be used as for diagnostic or other medical purposes.  
St John's Laboratory Ltd, Knowledge Dock Business Centre, University Way, London, E16 2RD | Tel: 0208 223 3081