

**Anti-BAG3 antibody (380-575) (STJ117026)**  
STJ117026

**GENERAL INFORMATION**


<b>Product Type</b>	Primary antibodies
<b>Short Description</b>	
<b>Applications</b>	WB/IHC-P/ELISA
<b>Host/Source</b>	Rabbit
<b>Reactivity</b>	Human/Mouse/Rat

**PRODUCT PROPERTIES**


<b>Clonality</b>	Polyclonal
<b>Clone ID</b>	
<b>Concentration</b>	Lot specific
<b>Conjugation</b>	Unconjugated
<b>Purification</b>	Affinity purification
<b>Dilution</b>	WB:1:500-1:2000
<b>Range</b>	IHC-P:1:50-1:100
	ELISA:Recommended starting concentration is 1 Mu g/mL. Please optimize the concentration based on your specific assay requirements.
<b>Formulation</b>	PBS with 0.01% Thimerosal, 50% Glycerol, pH 7.3.
<b>Isotype</b>	IgG
<b>Storage Instruction</b>	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

**TARGET INFORMATION**


<b>Gene ID</b>	9531
<b>Gene Symbol</b>	BAG3
<b>Uniprot ID</b>	BAG3_HUMAN
<b>Immunogen</b>	
<b>Immunogen Region</b>	380-575
<b>Specificity</b>	Recombinant fusion protein containing a sequence corresponding to amino acids 380-575 of human BAG3 (NP_004272.2).
<b>Immunogen Sequence</b>	PSAVPSSPKSVATEERAAPS TAPAEATPPKPGAEAPPKH PGVLKVEAILEKVQGLEQAV DNFEGKKTDKKYLMIEEYLT KELLALDSVDPEGRADVRQA RRDGVRKVQTILEKLEQKAI DVPGVQVQVYELQPSNLEADQ PLQAIMEMGAVAADKGGKNA GNAEDPHTTETQQPEATAAAT SNPSSMTDTPGNPAAP




Immunohistochemistry analysis of BAG3 in paraffin-embedded mouse heart using BAG3 Rabbit polyclonal antibody (STJ117026) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with immunohistochemistry staining protocol.



Immunohistochemistry analysis of BAG3 in paraffin-embedded human lung using BAG3 Rabbit polyclonal antibody (STJ117026) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with immunohistochemistry staining protocol.



Immunohistochemistry analysis of BAG3 in paraffin-embedded rat heart using BAG3 Rabbit polyclonal antibody (STJ117026) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with immunohistochemistry staining protocol.



Western blot analysis of various lysates using BAG3 Rabbit polyclonal antibody (STJ117026) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25 Mu g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 30s.

This product is suitable for in-vitro studies under the RESEARCH USE ONLY [RUO] licence. This product must not be used as for diagnostic or other medical purposes.  
St John's Laboratory Ltd, Knowledge Dock Business Centre, University Way, London, E16 2RD | Tel: 0208 223 3081