

Anti-OXCT2 antibody (430-510) (STJ117119)

STJ117119

GENERAL INFORMATION

Product Type Primary antibodies

Short Description

Applications WB/IF/ICC/ELISA Host/Source Rabbit

Reactivity Human/Mouse/Rat

PRODUCT PROPERTIES

Clonality Polyclonal

Clone ID

Concentration Lot specific
Conjugation Unconjugated
Purification Affinity purification
Dilution Range WB:1:500-1:2000

IF/ICC:1:50-1:200

ELISA:Recommended starting concentration is 1 Mu g/mL. Please optimize the concentration based on your specific assay

requirements.

Formulation PBS with 0.01% Thimerosal, 50% Glycerol, pH 7.3.

Isotype IgG

Storage Instruction Store at-20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

TARGET INFORMATION

Gene ID 64064 Gene Symbol OXCT2

Uniprot ID SCOT2_HUMAN


Immunogen

Immunogen Region 430-510


Specificity Recombinant fusion protein containing a sequence corresponding to amino acids 430-510 of human OXCT2 (NP_071403.1).

Immunogen QKTRVVVTMQHCTKDNTPKI MEKCTMPLTGKRCVDRIITE KAVFDVHRKKELTLRELWEG LTVDDIKKSTGCAFAVSPNL R


Immunogen Sequence



Immunofluorescence analysis of paraffin-embedde mouse testis using OXCT2 Rabbit polyclonal antibod (STJ117119) at dilution of 1:100. Secondary antibod, Cy3 Goat Anti-Rabbit IgG (H+L) at 1:500 dilution. Blue DAPI for nuclear staining.



Immunofluorescence analysis of paraffin-embedded ratestis using OXCT2 Rabbit polyclonal antibod (STJ117119) at dilution of 1:100. Secondary antibod Cy3 Goat Anti-Rabbit IgG (H+L) at 1:500 dilution. Blue DAPI for nuclear staining.



Western blot analysis of various lysates using OXCT2 Rabbit polyclonal antibody (STJ117119) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit (gG (H+L) (STJS000856) at 1:10000 dilution. Lysates/protiens: 25 Mu g per lane. Blocking buffer 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 5s