

Anti-SEPT8 antibody (348-427) (STJ117206)

STJ117206

GENERAL INFORMATION


Product Type	Primary antibodies
Short Description	
Applications	WB/IHC-P/IF/ICC/ELISA
Host/Source	Rabbit
Reactivity	Human/Mouse/Rat

PRODUCT PROPERTIES


Clonality	Polyclonal
Clone ID	
Concentration	Lot specific
Conjugation	Unconjugated
Purification	Affinity purification
Dilution Range	WB:1:500-1:1000 IHC-P:1:50-1:200 IF/ICC:1:50-1:200 ELISA:Recommended starting concentration is 1 Mu g/mL. Please optimize the concentration based on your specific assay requirements.
Formulation	PBS with 0.01% Thimerosal, 50% Glycerol, pH 7.3.
Isotype	IgG
Storage Instruction	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

TARGET INFORMATION


Gene ID	23176
Gene Symbol	SEPTIN8
Uniprot ID	SEPT8_HUMAN
Immunogen	
Immunogen Region	348-427
Specificity	Recombinant fusion protein containing a sequence corresponding to amino acids 348-427 of human Septin 8 (NP_001287728.1).
Immunogen Sequence	NKVKETELKEKERELHEK FEHLKRVHQEEKRKVEEKRR ELEETNAFNRRKA AVEALQ SQALHATSQQPLRKDKDKKN




Immunofluorescence analysis of U-2 OS cells using Septin 8 Rabbit polyclonal antibody (STJ117206) at dilution of 1:100. Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of L929 cells using Septin 8 Rabbit polyclonal antibody (STJ117206) at dilution of 1:100. Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of C6 cells using Septin 8 Rabbit polyclonal antibody (STJ117206) at dilution of 1:100. Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunohistochemistry analysis of Septin 8 in paraffin-embedded Human thyroid cancer using Septin 8 Rabbit polyclonal antibody (STJ117206) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with immunohistochemistry staining protocol.

This product is suitable for in-vitro studies under the RESEARCH USE ONLY [RUO] licence. This product must not be used as for diagnostic or other medical purposes.
St John's Laboratory Ltd, Knowledge Dock Business Centre, University Way, London, E16 2RD | Tel: 0208 223 3081