

**Anti-F10 antibody (41-300) (STJ117251)**  
STJ117251

**GENERAL INFORMATION**


|                          |                    |
|--------------------------|--------------------|
| <b>Product Type</b>      | Primary antibodies |
| <b>Short Description</b> |                    |
| <b>Applications</b>      | WB/IF/ICC/ELISA    |
| <b>Host/Source</b>       | Rabbit             |
| <b>Reactivity</b>        | Human/Mouse/Rat    |

**PRODUCT PROPERTIES**


|                            |                                                                                                                                                            |
|----------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Clonality</b>           | Polyclonal                                                                                                                                                 |
| <b>Clone ID</b>            |                                                                                                                                                            |
| <b>Concentration</b>       | Lot specific                                                                                                                                               |
| <b>Conjugation</b>         | Unconjugated                                                                                                                                               |
| <b>Purification</b>        | Affinity purification                                                                                                                                      |
| <b>Dilution</b>            | WB:1:500-1:2000                                                                                                                                            |
| <b>Range</b>               | IF/CC:1:50-1:200<br>ELISA:Recommended starting concentration is 1 $\mu$ g/mL. Please optimize the concentration based on your specific assay requirements. |
| <b>Formulation</b>         | PBS with 0.01% Thimerosal, 50% Glycerol, pH 7.3.                                                                                                           |
| <b>Isotype</b>             | IgG                                                                                                                                                        |
| <b>Storage Instruction</b> | Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.                                                             |

**TARGET INFORMATION**


|                           |                                                                                                                                                                                                                                                                  |
|---------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Gene ID</b>            | 2159                                                                                                                                                                                                                                                             |
| <b>Gene Symbol</b>        | F10                                                                                                                                                                                                                                                              |
| <b>Uniprot ID</b>         | FA10_HUMAN                                                                                                                                                                                                                                                       |
| <b>Immunogen</b>          |                                                                                                                                                                                                                                                                  |
| <b>Immunogen Region</b>   | 41-300                                                                                                                                                                                                                                                           |
| <b>Specificity</b>        | Recombinant fusion protein containing a sequence corresponding to amino acids 41-300 of human F10 (NP_000495.1).                                                                                                                                                 |
| <b>Immunogen Sequence</b> | ANSFLEEMKKGHLERECMEE TCSYEEAREVFEDSDKTNEF WNKYKDGDCQETSPCQNQGK CKDGLGEYTCTCLEGFEGKN<br>CELFTRKLCSLDNGDCDQFC HEEQNSVVCSCARGYTLADN GKACIPTGPYPGKGQTLERR KRSVAQATSSSGEAPDSITW<br>KPYDAADLDPTENPFDLLDF NQTQPERGDNNLTRIVGGQE CKDGCEPWQALLINEENEGF CGGTILSEFYILTAACHLY |



Immunofluorescence analysis of paraffin-embedded mouse liver using F10 Rabbit polyclonal antibody (STJ117251) at dilution of 1:100 (40x lens). Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of paraffin-embedded rat liver using F10 Rabbit polyclonal antibody (STJ117251) at dilution of 1:100 (40x lens). Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) at 1:500 dilution. Blue: DAPI for nuclear staining.



Western blot analysis of various lysates using F10 Rabbit polyclonal antibody (STJ117251) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJ500856) at 1:10000 dilution. Lysates/proteins: 25  $\mu$ g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 10s.

This product is suitable for in-vitro studies under the RESEARCH USE ONLY [RUO] licence. This product must not be used as for diagnostic or other medical purposes.  
St John's Laboratory Ltd, Knowledge Dock Business Centre, University Way, London, E16 2RD | Tel: 0208 223 3081