

Anti-SLC17A6 antibody (500-582) (STJ117371)

STJ117371

GENERAL INFORMATION


Product Type	Primary antibodies
Short Description	
Applications	WB/IF/ICC/ELISA
Host/Source	Rabbit
Reactivity	Human/Mouse/Rat

PRODUCT PROPERTIES


Clonality	Polyclonal
Clone ID	
Concentration	Lot specific
Conjugation	Unconjugated
Purification	Affinity purification
Dilution Range	WB:1:500-1:2000 IF/ICC:1:50-1:200 ELISA:Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.
Formulation	PBS with 0.05% Proclin300, 50% Glycerol, pH 7.3.
Isotype	IgG
Storage Instruction	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

TARGET INFORMATION


Gene ID	57084
Gene Symbol	SLC17A6
Uniprot ID	VGLU2_HUMAN
Immunogen	
Immunogen Region	500-582
Specificity	A synthetic peptide corresponding to a sequence within amino acids 500-582 of human SLC17A6 (NP_065079.1).
Immunogen Sequence	GEKQPWADPEETSEKCGFI HEDELDEETGDITQNYINYG TTKSYGATTQANGGWPSGWEE KKEEFVQGEVQDHSYKDRV DYS



Immunofluorescence analysis of mouse brain using SLC17A6 Rabbit polyclonal antibody (STJ117371) at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.



Immunofluorescence analysis of rat brain using SLC17A6 Rabbit polyclonal antibody (STJ117371) at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.



Western blot analysis of extracts of various cell lines, using SLC17A6 Polyclonal Antibody (STJ117371) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 10s.