

Anti-ATG12 antibody (1-100) (STJ118058) STJ118058

GENERAL INFORMATION

Product Type Primary antibodies Short Description Applications WB/IF/ICC/ELISA Host/Source Rabbit Reactivity Human/Mouse/Rat

PRODUCT PROPERTIES

 Clonality
 Polyclonal

 Clone ID
 Concentration

 Concentration
 Lot specific

 Concinguation
 Unconjugated

 Purification
 Affinity purification

 Plutition Range
 WB:1:500-1:1000


 IF/CC:1:50-1:200
 ELISA:Recommended starting concentration is 1 Mu g/mL. Please optimize the concentration based on your specific assay requirements.

 Formulation
 PBS with 0.05% Proclin300, 50% Glycerol, pH 7.3.


 Isotype
 IgG

 Storage Instruction
 Stor at-20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.


TARGET INFORMATION




Immunofluorescence analysis of U2OS cells using ATG12 Rabbit polyclonal antibody (STJ118058) at dilution of 1:50 (40x lens). Blue: DAPI for nuclear staining.



Immunofluorescence analysis of PC-12 cells using ATG12 Rabbit polyclonal antibody (STJ118058) at dilution of 1:50 (40x lens). Blue: DAPI for nuclear stainion



Immunofluorescence analysis of NIH/3T3 cells using ATG12 Rabbit polyclonal antibody (STJ118058) at dilution of 1:50 (40x lens). Blue: DAPI for nuclear staining.



Western blot analysis of extracts of various cell lines, using ATG12 antibody (STJ118058) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H-L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25 Mu g per lane. Blocking buffer: 3% nontrat dry milk in TBS1. Detection: ECL Enhanced Kit. Exposure time: 1900.

This product is suitable for in-vitro studies under the RESEARCH USE ONLY [RUO] licence. This product must not be used as for diagnostic or other medical purposes. St John's Laboratory Ltd, Knowledge Dock Business Centre, University Way, London, E16 2RD | Tel: 0208 223 3081