

Anti-DCT antibody (50-350) (STJ118481)

STJ118481

GENERAL INFORMATION


Product Type	Primary antibodies
Short Description	
Applications	WB/ELISA
Host/Source	Rabbit
Reactivity	Human/Mouse

PRODUCT PROPERTIES

Clonality	Polyclonal
Clone ID	
Concentration	Lot specific
Conjugation	Unconjugated
Purification	Affinity purification
Dilution	WB:1:500-1:2000
Range	ELISA:Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.
Formulation	PBS with 0.01% Thimerosal, 50% Glycerol, pH 7.3.
Isotype	IgG
Storage	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.
Instruction	

TARGET INFORMATION

Gene ID	1638
Gene Symbol	DCT
Uniprot ID	TYRP2_HUMAN
Immunogen	
Immunogen Region	50-350
Specificity	Recombinant fusion protein containing a sequence corresponding to amino acids 50-350 of human DCT (NP_001913.2).
Immunogen Sequence	NVCGSQQGRGQCTEVRADTR PWSGPYILRNQDDRELWPRK FFHRTCKCTGNFAGYNCGDC KFGWTGPNCERKKPPVIRQN IHSLSPQEREQFLGALDLAK KRVHPDYVITQHWLGLLGP NGTQPFANCNVYDFVWLH YYSVRDTLLGPGRPYRAIDF SHQGPAFVTWHRYHLLCLER DLQRLIGNESFALPYWNFAT GRNECDVCTDQLFGAARPDD PTLISRNSRFSSWETVCD



Western blot analysis of various lysates using DCT Rabbit polyclonal antibody (STJ118481) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 10s.

This product is suitable for in-vitro studies under the RESEARCH USE ONLY [RUO] licence. This product must not be used as for diagnostic or other medical purposes.
St John's Laboratory Ltd, Knowledge Dock Business Centre, University Way, London, E16 2RD | Tel: 0208 223 3081