

Anti-STAP2 antibody (130-220) (STJ118571)

STJ118571

GENERAL INFORMATION


Product Type	Primary antibodies
Short Description	
Applications	WB/ELISA
Host/Source	Rabbit
Reactivity	Human/Mouse

PRODUCT PROPERTIES


Clonality	Polyclonal
Clone ID	
Concentration	Lot specific
Conjugation	Unconjugated
Purification	Affinity purification
Dilution Range	WB:1:500-1:2000 ELISA:Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.
Formulation	PBS with 0.01% Thimerosal, 50% Glycerol, pH 7.3.
Isotype	IgG
Storage Instruction	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

TARGET INFORMATION

Gene ID	55620
Gene Symbol	STAP2
Uniprot ID	STAP2_HUMAN
Immunogen	
Immunogen Region	130-220
Specificity	Recombinant fusion protein containing a sequence corresponding to amino acids 130-220 of human STAP2 (NP_001013863).
Immunogen Sequence	GHLYMMSEVLAKKEARRALE TPSCFLKVSRLAEQLLLERY PECGNLLLRPSGDGADGVSV TTRQMHNGTHVVRHYKVKRE GPKYVIDVEQP



Western blot analysis of lysates from HT-29 cells, using STAP2 Rabbit polyclonal antibody (STJ118571) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 240s.



Western blot analysis of lysates from Mouse heart, using STAP2 Rabbit polyclonal antibody (STJ118571) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 90s.

This product is suitable for in-vitro studies under the RESEARCH USE ONLY [RUO] licence. This product must not be used as for diagnostic or other medical purposes.
St John's Laboratory Ltd, Knowledge Dock Business Centre, University Way, London, E16 2RD | Tel: 0208 223 3081