

Anti-FKBP1B antibody (1-80) (STJ118803)

STJ118803

GENERAL INFORMATION


Product Type	Primary antibodies
Short Description	
Applications	WB/IHC-P/IF/ICC/ELISA
Host/Source	Rabbit
Reactivity	Human/Mouse/Rat

PRODUCT PROPERTIES


Clonality	Polyclonal
Clone ID	
Concentration	Lot specific
Conjugation	Unconjugated
Purification	Affinity purification
Dilution Range	WB:1:500-1:2000 IHC-P:1:50-1:100 IF/ICC:1:50-1:100 ELISA:Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.
Formulation	PBS with 0.01% Thimerosal, 50% Glycerol, pH 7.3.
Isotype	IgG
Storage Instruction	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

TARGET INFORMATION


Gene ID	2281
Gene Symbol	FKBP1B
Uniprot ID	FKB1B_HUMAN
Immunogen	
Immunogen Region	1-80
Specificity	A synthetic peptide corresponding to a sequence within amino acids 1-80 of human FKBP1B (NP_473374.1).
Immunogen Sequence	MGVEIETISPGDGRTPKKG QTCVVHYTGMLQNGKKFDSS RDRNKPFRIGKQEVKGF EEGAAQLGPLSLPICPHPC




Immunofluorescence analysis of L929 cells using FKBP1B Rabbit polyclonal antibody (STJ118803) at dilution of 1:100 (40x lens). Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunohistochemistry analysis of FKBP1B in paraffin-embedded mouse heart using FKBP1B Rabbit polyclonal antibody (STJ118803) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with immunohistochemistry staining protocol.



Immunohistochemistry analysis of FKBP1B in paraffin-embedded mouse spinal cord using FKBP1B Rabbit polyclonal antibody (STJ118803) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with immunohistochemistry staining protocol.



Immunohistochemistry analysis of FKBP1B in paraffin-embedded rat heart using FKBP1B Rabbit polyclonal antibody (STJ118803) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with immunohistochemistry staining protocol.

This product is suitable for in-vitro studies under the RESEARCH USE ONLY [RUO] licence. This product must not be used as for diagnostic or other medical purposes.
St John's Laboratory Ltd, Knowledge Dock Business Centre, University Way, London, E16 2RD | Tel: 0208 223 3081