

Anti-UGT2B15 antibody (60-190) (STJ119076)

STJ119076

GENERAL INFORMATION


Product Type	Primary antibodies
Short Description	
Applications	WB/IHC-P/ELISA
Host/Source	Rabbit
Reactivity	Human/Mouse/Rat

PRODUCT PROPERTIES


Clonality	Polyclonal
Clone ID	
Concentration	Lot specific
Conjugation	Unconjugated
Purification	Affinity purification
Dilution Range	WB:1:500-1:2000 IHC-P:1:100-1:200 ELISA:Recommended starting concentration is 1 Mu g/mL. Please optimize the concentration based on your specific assay requirements.
Formulation	PBS with 0.01% Thimerosal, 50% Glycerol, pH 7.3.
Isotype	IgG
Storage Instruction	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

TARGET INFORMATION


Gene ID	7366
Gene Symbol	UGT2B15
Uniprot ID	UDB15_HUMAN
Immunogen	
Immunogen Region	60-190
Specificity	Recombinant fusion protein containing a sequence corresponding to amino acids 60-190 of human UGT2B15 (NP_001067.2).
Immunogen Sequence	ASTLVNASKSSAIKLEVYPT SLTKNYLEDSSLKILDRWIY GVSKNTFWSYFSQLQELCWE YYDYSNKLCKDAVLNKKLMM KLQESKFDVILADALNPCGE LLAELFNIPFLYSLRFSVG Y TFEKNGGGFLF




Immunohistochemistry analysis of UGT2B15 in paraffin-embedded mouse spleen using UGT2B15 Rabbit polyclonal antibody (STJ119076) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with immunohistochemistry staining protocol.



Immunohistochemistry analysis of UGT2B15 in paraffin-embedded human oophoroma using UGT2B15 Rabbit polyclonal antibody (STJ119076) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with immunohistochemistry staining protocol.



Immunohistochemistry analysis of UGT2B15 in paraffin-embedded rat kidney using UGT2B15 Rabbit polyclonal antibody (STJ119076) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with immunohistochemistry staining protocol.



Western blot analysis of various lysates using UGT2B15 Rabbit polyclonal antibody (STJ119076) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25 Mu g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 5s.

This product is suitable for in-vitro studies under the RESEARCH USE ONLY [RUO] licence. This product must not be used as for diagnostic or other medical purposes.
St John's Laboratory Ltd, Knowledge Dock Business Centre, University Way, London, E16 2RD | Tel: 0208 223 3081