

Anti-PDCD10 antibody (1-212) (STJ119307)
STJ119307

GENERAL INFORMATION



Product Type	Primary antibodies
Short Description	
Applications	WB/IHC-P/IF/ICC/ELISA
Host/Source	Rabbit
Reactivity	Human/Mouse/Rat

PRODUCT PROPERTIES


Clonality	Polyclonal
Clone ID	
Concentration	Lot specific
Conjugation	Unconjugated
Purification	Affinity purification
Dilution	WB:1:500-1:1000
Range	IHC-P:1:50-1:200 IF/ICC:1:50-1:200 ELISA:Recommended starting concentration is 1 μ g/mL. Please optimize the concentration based on your specific assay requirements.
Formulation	PBS with 0.05% Proclin300, 50% Glycerol, pH 7.3.
Isotype	IgG
Storage	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.
Instruction	

TARGET INFORMATION


Gene ID	11235
Gene Symbol	PDCD10
Uniprot ID	PDC10_HUMAN
Immunogen	
Immunogen Region	1-212
Specificity	Recombinant fusion protein containing a sequence corresponding to amino acids 1-212 of human PDCD10 (NP_009148.2).
Immunogen Sequence	MRMTMEEMKNEAETSMVSM PLYAVMYPVFNELERVNLSA AQTLRAAFIKAENPGLTQ DIIMKILEKKSVEVNFTESL LRMAADDVEEYMIERPEPEF QDLNEKARALKQILSKIPDE INDRVRLFQTIKDIAIAIKE LLDTVNNVFKKYQYNRRAL EHQKKEFVKYSKFSFDLTKT YFKDGGKAINVFSANRLIHQ TNLILQFTKTVA



Immunohistochemistry analysis of paraffin-embedded rat spleen using PDCD10 antibody (STJ119307) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7. 2 before commencing with immunohistochemistry staining protocol.



Immunohistochemistry analysis of paraffin-embedded human breast cancer using PDCD10 antibody (STJ119307) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7. 2 before commencing with immunohistochemistry staining protocol.



Immunohistochemistry analysis of paraffin-embedded mouse kidney using PDCD10 antibody (STJ119307) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7. 2 before commencing with immunohistochemistry staining protocol.

This product is suitable for in-vitro studies under the RESEARCH USE ONLY [RUO] licence. This product must not be used as for diagnostic or other medical purposes.
St John's Laboratory Ltd, Knowledge Dock Business Centre, University Way, London, E16 2RD | Tel: 0208 223 3081