

Anti-FCF1 antibody (100-198) (STJ119324)

STJ119324

GENERAL INFORMATION


Product Type	Primary antibodies
Short Description	
Applications	WB/IHC-P/IF/ICC/ELISA
Host/Source	Rabbit
Reactivity	Human/Rat

PRODUCT PROPERTIES


Clonality	Polyclonal
Clone ID	
Concentration	Lot specific
Conjugation	Unconjugated
Purification	Affinity purification
Dilution Range	WB:1:500-1:2000 IHC-P:1:50-1:200 IF/ICC:1:50-1:200 ELISA:Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.
Formulation	PBS with 0.01% Thimerosal, 50% Glycerol, pH 7.3.
Isotype	IgG
Storage Instruction	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

TARGET INFORMATION


Gene ID	51077
Gene Symbol	FCF1
Uniprot ID	FCF1_HUMAN
Immunogen	
Immunogen Region	100-198
Specificity	A synthetic peptide corresponding to a sequence within amino acids 100-198 of human FCF1 (NP_057046.1).
Immunogen Sequence	PCITDCVMAIEIKLGQKYRV ALRIAKDPRFERLPCTHKGT YADDCLVQRVTQHKCYIVAT VDRDLKRRIRKIPGVPMYI SNHRYNIERPDPDYGAPRF




Immunofluorescence analysis of U-2 OS cells using FCF1 Rabbit polyclonal antibody (STJ119324) at dilution of 1:100. Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of C6 cells using FCF1 Rabbit polyclonal antibody (STJ119324) at dilution of 1:100. Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunohistochemistry analysis of FCF1 in paraffin-embedded Human esophageal using FCF1 Rabbit polyclonal antibody (STJ119324) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with immunohistochemistry staining protocol.



Western blot analysis of lysates from 293T cells, using FCF1 Rabbit polyclonal antibody (STJ119324) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 100s.

This product is suitable for in-vitro studies under the RESEARCH USE ONLY [RUO] licence. This product must not be used as for diagnostic or other medical purposes.

St John's Laboratory Ltd, Knowledge Dock Business Centre, University Way, London, E16 2RD | Tel: 0208 223 3081