

## Anti-FCGR1B antibody (100-200) (STJ119586)

STJ119586

### GENERAL INFORMATION


<b>Product Type</b>	Primary antibodies
<b>Short Description</b>	
<b>Applications</b>	WB/IHC-P/ELISA
<b>Host/Source</b>	Rabbit
<b>Reactivity</b>	Human

### PRODUCT PROPERTIES


<b>Clonality</b>	Polyclonal
<b>Clone ID</b>	
<b>Concentration</b>	Lot specific
<b>Conjugation</b>	Unconjugated
<b>Purification</b>	Affinity purification
<b>Dilution Range</b>	WB:1:500-1:2000 IHC-P:1:50-1:200 ELISA:Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.
<b>Formulation</b>	PBS with 0.01% Thimerosal, 50% Glycerol, pH 7.3.
<b>Isotype</b>	IgG
<b>Storage</b>	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.
<b>Instruction</b>	

### TARGET INFORMATION


<b>Gene ID</b>	
<b>Gene Symbol</b>	FCGR1BP
<b>Uniprot ID</b>	FCGRB_HUMAN
<b>Immunogen</b>	
<b>Immunogen Region</b>	100-200
<b>Specificity</b>	A synthetic peptide corresponding to a sequence within amino acids 100-200 of human FCGR1B (NP_001017986.1).
<b>Immunogen Sequence</b>	IHRGWLLQVSSRVFMEGEP LALRCHAWKDKLVYNVLYYR NGKAFKFFHWNSNLTKTN ISHNGTYHCSGMGKHRYTSA GISQYTVKGLQLPTPVWFHV L



Immunohistochemistry analysis of FCGR1B in paraffin-embedded human placenta using FCGR1B Rabbit polyclonal antibody (STJ119586) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with immunohistochemistry staining protocol.



Immunohistochemistry analysis of FCGR1B in paraffin-embedded human mammary cancer using FCGR1B Rabbit polyclonal antibody (STJ119586) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with immunohistochemistry staining protocol.



Immunohistochemistry analysis of FCGR1B in paraffin-embedded human lung cancer using FCGR1B Rabbit polyclonal antibody (STJ119586) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with immunohistochemistry staining protocol.