

Anti-LYAR antibody (10-160) (STJ119766)

STJ119766

GENERAL INFORMATION


Product Type	Primary antibodies
Short Description	
Applications	WB/IHC-P/IF/ICC/ELISA
Host/Source	Rabbit
Reactivity	Human/Mouse/Rat

PRODUCT PROPERTIES


Clonality	Polyclonal
Clone ID	
Concentration	Lot specific
Conjugation	Unconjugated
Purification	Affinity purification
Dilution	WB:1:500-1:2000
Range	IHC-P:1:50-1:200 IF/ICC:1:50-1:200 ELISA:Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.
Formulation	PBS with 0.01% Thimerosal, 50% Glycerol, pH 7.3.
Isotype	IgG
Storage	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.
Instruction	

TARGET INFORMATION


Gene ID	55646
Gene Symbol	LYAR
Uniprot ID	LYAR_HUMAN
Immunogen	
Region	10-160
Specificity	Recombinant fusion protein containing a sequence corresponding to amino acids 10-160 of human LYAR (NP_060286.1).
Immunogen Sequence	GESVKKIQVEKHVSVCNCE CLSCIDCGKDFWGDDYKNHV KCISEDQKYGGKGYEGKTHK GDIKQQAIIQKISELIKRPN VSPKVRLELQISAFDNVPR KKAQFQNMWKNLKVHNESI LDQVWNIFSEASNSEPVNKE QDQRPLHPVAN




Immunofluorescence analysis of C6 cells using LYAR Rabbit polyclonal antibody (STJ119766) at dilution of 1:100. Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of L929 cells using LYAR Rabbit polyclonal antibody (STJ119766) at dilution of 1:100. Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of U2OS cells using LYAR Rabbit polyclonal antibody (STJ119766) at dilution of 1:100. Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunohistochemistry analysis of LYAR in paraffin-embedded rat testis using LYAR Rabbit polyclonal antibody (STJ119766) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with immunohistochemistry staining protocol.

This product is suitable for in-vitro studies under the RESEARCH USE ONLY [RUO] licence. This product must not be used as for diagnostic or other medical purposes.
St John's Laboratory Ltd, Knowledge Dock Business Centre, University Way, London, E16 2RD | Tel: 0208 223 3081