

## Anti-CAMK2A antibody (300-400) (STJ119903)

STJ119903

### GENERAL INFORMATION


<b>Product Type</b>	Primary antibodies
<b>Short Description</b>	
<b>Applications</b>	WB/IF/ICC/ELISA
<b>Host/Source</b>	Rabbit
<b>Reactivity</b>	Human/Mouse/Rat

### PRODUCT PROPERTIES


<b>Clonality</b>	Polyclonal
<b>Clone ID</b>	
<b>Concentration</b>	Lot specific
<b>Conjugation</b>	Unconjugated
<b>Purification</b>	Affinity purification
<b>Dilution Range</b>	WB:1:500-1:1000 IF/ICC:1:50-1:200 ELISA:Recommended starting concentration is 1 Mu g/mL. Please optimize the concentration based on your specific assay requirements.
<b>Formulation</b>	PBS with 0.05% Proclin300, 50% Glycerol, pH 7.3.
<b>Isotype</b>	IgG
<b>Storage Instruction</b>	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

### TARGET INFORMATION

<b>Gene ID</b>	815
<b>Gene Symbol</b>	CAMK2A
<b>Uniprot ID</b>	KCC2A_HUMAN
<b>Immunogen</b>	
<b>Immunogen Region</b>	300-400
<b>Specificity</b>	A synthetic peptide corresponding to a sequence within amino acids 300-400 of human CAMK2A (NP_741960.1).
<b>Immunogen Sequence</b>	KGAILTTMLATRNFGGKSG GNKKSDGVKESSESTNTTIE DEDTKVRKQEIIKVTEQLIE AISNGDFESYTKMCDPGMTA FEPEALGNLVEGLDFHRFYF E



Immunofluorescence analysis of NIH/3T3 cells using CAMK2A Rabbit polyclonal antibody (STJ119903) at dilution of 1:100. Blue: DAPI for nuclear staining.



Western blot analysis of various lysates using CAMK2A Rabbit polyclonal antibody (STJ119903) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25 Mu g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 180s.

This product is suitable for in-vitro studies under the RESEARCH USE ONLY [RUO] licence. This product must not be used as for diagnostic or other medical purposes.  
St John's Laboratory Ltd, Knowledge Dock Business Centre, University Way, London, E16 2RD | Tel: 0208 223 3081