

Anti-MT-CO2 antibody (40-100) (STJ119944)

STJ119944

GENERAL INFORMATION



Product Type	Primary antibodies
Short Description	
Applications	WB/IHC-P/IF/ICC/ELISA
Host/Source	Rabbit
Reactivity	Human/Mouse/Rat

PRODUCT PROPERTIES


Clonality	Polyclonal
Clone ID	
Concentration	Lot specific
Conjugation	Unconjugated
Purification	Affinity purification
Dilution Range	WB:1:1000-1:5000 IHC-P:1:50-1:200 IF/ICC:1:50-1:200 ELISA:Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.
Formulation	PBS with 0.05% Proclin300, 50% Glycerol, pH 7.3.
Isotype	IgG
Storage Instruction	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

TARGET INFORMATION


Gene ID	4513
Gene Symbol	MT-CO2
Uniprot ID	COX2_HUMAN
Immunogen	
Immunogen Region	40-100
Specificity	A synthetic peptide corresponding to a sequence within amino acids 40-100 of human MTCO2 (YP_003024029.1).
Immunogen Sequence	YALFLTLTKLTNTNISDAQ EMETVWTILPAIILVLIALP SLRILYMTDEVNDPSLTIKS I



Immunohistochemistry analysis of paraffin-embedded human colon using MTCO2 rabbit polyclonal antibody (STJ119944) at dilution of 1:100 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with immunohistochemistry staining protocol.



Immunohistochemistry analysis of paraffin-embedded rat colon using MTCO2 rabbit polyclonal antibody (STJ119944) at dilution of 1:100 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with immunohistochemistry staining protocol.



Immunofluorescence analysis of HepG2 cells using MTCO2 rabbit polyclonal antibody (STJ119944) at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.