

Anti-MT-ND2 antibody (248-347) (STJ119947)

STJ119947

GENERAL INFORMATION


Product Type	Primary antibodies
Short Description	WB/IHC-P/IF/ICC/ELISA
Applications	Rabbit
Host/Source	Human/Mouse/Rat
Reactivity	

PRODUCT PROPERTIES


Clonality	Polyclonal
Clone ID	
Concentration	Lot specific
Conjugation	Unconjugated
Purification	Affinity purification
Dilution Range	WB:1:500-1:1000 IHC-P:1:50-1:200 IF/ICC:1:50-1:200
	ELISA:Recommended starting concentration is 1 Mu g/mL. Please optimize the concentration based on your specific assay requirements.
Formulation	PBS with 0.09% Sodium Azide, 50% Glycerol, pH 7.3.
Isotype	IgG
Storage Instruction	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

TARGET INFORMATION


Gene ID	4536
Gene Symbol	MT-ND2
Uniprot ID	NU2M_HUMAN
Immunogen	
Immunogen Region	248-347
Specificity	A synthetic peptide corresponding to a sequence within amino acids 248-347 of human MT-ND2 (YP_003024027.1).
Immunogen Sequence	LLSLGGLPPLTGFLPKWAI EEFTKNNNSLIPTIMATITL LNLYFYLRLIYSTSITLLPM SNNVKMKWQFEHTKPTPFLP TLIALTLLLPISPFLMIL



Immunofluorescence analysis of HeLa cells using MT-ND2 Rabbit polyclonal antibody (STJ119947) at dilution of 1:100. Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunohistochemistry analysis of MT-ND2 in paraffin-embedded human colon carcinoma using MT-ND2 Rabbit polyclonal antibody (STJ119947) at dilution of 1:50 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with immunohistochemistry staining protocol.



Western blot analysis of various lysates using MT-ND2 Rabbit polyclonal antibody (STJ119947) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJS000856) at 1:10000 dilution. Lanes/proteins: 25 Mu g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 90s.