

Anti-CD63 antibody (120aa to 175aa) (STJ140029)

GENERAL INFORMATION

Product Type Primary antibodies

Short Goat polyclonal antibody anti-Lysosome-associated membrane glycoprotein 3 (120aa to 175aa) is suitable for use in Western Blot,

Description Immunohistochemistry and Immunofluorescence research applications.

Host/Source Goat

Reactivity Human/Rat/Mouse/Monkey/Canine

PRODUCT PROPERTIES

Clonality Polyclonal

Clone ID Concentration 3 mg/mL Conjugation Unconjugated

Purification This antibody is epitope-affinity purified from goat antiserum.

Dilution WB 1:500-1:5000 Range IF 1:25-1:250 IHC-F 1:250-1:1000 IHC-P 1:250-1:1000

Amorim M Martins B Caramelo F et al. Front Med-Lausanne 2022 May PMID: 35692536

Matsui T Sakamaki Y Nakashima S et al. Cell Rep 2022 May PMID: 35649370

Martin

Formulation PBS, 20% Glycerol and 0.05% Sodium Azide.

Isotype IgG

Storage For continuous use, store at 2-8 C for one-two days. For extended storage, store in-20 C freezer. Working dilution samples should be

Instruction discarded if not used within 12 hours.

TARGET INFORMATION

Gene ID 27074

Gene Symbol LAMP3

Uniprot ID LAMP3_HUMAN

Immunogen Purified recombinant peptide derived from within residues 120 aa to 175 aa of human CD63 produced in E. coli.

Immunogen


Region

Specificity Reacts with CD63, a 40-60 kDa glycoprotein, detected by Western blot in the following human (HeLa, HUH, Jurkat) , mouse (AtT-20,


Hepa, 3T3, RAW264.7), canine (MDCK) and monkey (COS-7) whole cell lysates.

Immunogen MENYPKNNHTASILDRMQAD FKCCGAANYTDWEKIPSMSK NRVPDSCCI

Sequence



Anti-CD63 antibody at 1:2500 dilution lysate at 50 µg per lane Rabbit polyclonal to goat IgG (HRP) at 1:10000



Anti-CD63 antibody at 1:2500 dilution lysate at 50 µg per-lane Rabbit polyclonal to goat IgG (HRP) at 1:10000

