

Anti-CD57 antibody [NK-1] (STJ16100013)

GENERAL INFORMATION

Product Type Primary antibodies

Short Mouse monoclonal antibody anti-CD57 is suitable for use in Flow Cytometry, Immunofluorescence and Immunohistochemistry

Description research applications.

Applications FC/IF/IHC Host/Source Mouse Reactivity Human

PRODUCT PROPERTIES

Clonality Monoclonal Clone ID NK-1

 $\textbf{Concentration} \quad \text{Can be provided as 100 } \mu\text{g/mL}, 500 \; \mu\text{g/mL or 1mg/mL}.$

Conjugation Unconjugated

Purification Affinity purified from tissue culture.

 $\textbf{Dilution} \quad \text{Flow cytometry (0, 5-1, 0 μg/million cells in 0, 1 ml)}. \ Immunofluorescence (0, 5-1, 0 μg/ml). \ Immunohistology (1-2 μg/ml) for 30-60 min and 10 million cells in 0, 1 ml). \ Immunofluorescence (0, 5-1, 0 μg/ml). \ Immunohistology (1-2 μg/ml) for 30-60 min and 10 million cells in 0, 1 ml). \ Immunohistology (1-2 μg/ml) for 30-60 min and 10 million cells in 0, 1 ml). \ Immunohistology (1-2 μg/ml) for 30-60 min and 10 million cells in 0, 1 ml). \ Immunohistology (1-2 μg/ml) for 30-60 min and 10 million cells in 0, 1 ml). \ Immunohistology (1-2 μg/ml) for 30-60 min and 10 million cells in 0, 1 ml). \ Immunohistology (1-2 μg/ml) for 30-60 min and 10 million cells in 0, 1 ml). \ Immunohistology (1-2 μg/ml) for 30-60 min and 10 million cells in 0, 1 ml). \ Immunohistology (1-2 μg/ml) for 30-60 min and 10 million cells in 0, 1 ml). \ Immunohistology (1-2 μg/ml) for 30-60 min and 10 million cells in 0, 1 ml). \ Immunohistology (1-2 μg/ml) for 30-60 min and 10 million cells in 0, 1 ml). \ Immunohistology (1-2 μg/ml) for 30-60 min and 10 million cells in 0, 1 ml). \ Immunohistology (1-2 μg/ml) for 30-60 min and 10 million cells in 0, 1 ml). \ Immunohistology (1-2 μg/ml) for 30-60 min and 10 million cells in 0, 1 ml). \ Immunohistology (1-2 μg/ml) for 30-60 min and 10 million cells in 0, 1 ml). \ Immunohistology (1-2 μg/ml) for 30-60 min and 10 million cells in 0, 1 ml). \ Immunohistology (1-2 μg/ml) for 30-60 min and 10 million cells in 0, 1 ml). \ Immunohistology (1-2 μg/ml) for 30-60 min and 10 ml) for 30-60 min and 10 ml). \ Immunohistology (1-2 μg/ml) for 30-60 min and 10 ml) for 30-60$

Range at RT; staining of formalin-fixed tissues requires boiling tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for

Formulation PBS with 0.02% Sodium Azide.

Isotype IgMk

Storage Store for up to 1 year at 2-8°C upon receipt. **Instruction**

TARGET INFORMATION

Gene ID

Gene Symbol Uniprot ID

Immunogen A BALB/c mouse was immunized with human PBL. Fusion partner: NS-1.

Immunogen Region Specificity Immunogen Sequence

