

## Anti-Cytokeratin 20 antibody [ZR429] (STJ180643)

STJ180643

### GENERAL INFORMATION


|                          |   |
|--------------------------|---|
| <b>Product Type</b>      | Primary antibodies  |
| <b>Short Description</b> | Rabbit monoclonal antibody anti-Cytokeratin 20 is suitable for use in Immunohistochemistry research applications. |
| <b>Applications</b>      | IHC-P   |
| <b>Host/Source</b>       | Rabbit  |
| <b>Reactivity</b>        | Human   |

### PRODUCT PROPERTIES

|                       |  |
|-----------------------|--|
| <b>Clonality</b>      | Monoclonal   |
| <b>Clone ID</b>       | ZR429  |
| <b>Concentration</b>  |  |
| <b>Conjugation</b>    | Unconjugated   |
| <b>Purification</b>   | Affinity purified  |
| <b>Dilution Range</b> | 1:100-200  |
| <b>Formulation</b>    | Tris-HCl buffer containing stabilizing protein (BSA) and <0.1% ProClin   |
| <b>Isotype</b>        |  |
| <b>Storage</b>        | Store at 2-8°C for up to 24 months. Predilute: Ready to use, no reconstitution necessary. Concentrate: Use dilution range and appropriate lab-standardized diluent. Stability after dilution: 7 days at 24°C, 3 months at 2-8°C, 6months at -20°C. |
| <b>Instruction</b>    |  |

### TARGET INFORMATION

**Gene ID**  
**Gene Symbol**  
**Uniprot ID**  
**Immunogen**  
**Immunogen**  
**Region**  
**Specificity**  
**Immunogen**  
**Sequence**



Human colon carcinoma stained with anti-keratin 20 antibody using peroxidase-conjugate and DAB chromogen. Note the cytoplasmic staining of carcinoma glands.