

## Anti-Phospho-FOXO4-S197 antibody (STJ22147)

ST.122147

## **GENERAL INFORMATION**

Product Type Primary antibodies

**Short Description** 

Applications WB/IF/ICC/ELISA Host/Source Rabbit

Reactivity Human/Mouse/Rat

## **PRODUCT PROPERTIES**

Clonality Polyclonal

Clone ID

Concentration Lot specific
Conjugation Unconjugated
Purification Affinity purification
Dilution Range WB:1:100-1:500

IF/ICC:1:100-1:200

ELISA:Recommended starting concentration is 1 Mu g/mL. Please optimize the concentration based on your specific assay

requirements.

Formulation PBS with 0.05% Proclin300, 50% Glycerol, pH 7.3.

**Isotype** IgG

Storage Instruction Store at-20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

## **TARGET INFORMATION**

Gene ID 4303
Gene Symbol FOXO4
Uniprot ID FOXO4\_HUMAN


Immunogen AASMD

Immunogen Region

Specificity A synthetic phosphorylated peptide around S197 of human FOXO4 (NP\_005929.2).


Immunogen AASMD

Sequence





Western blot analysis of extracts from 293 cells using phospho-FOXO4-S197 antibody (STJ22147). Secondary antibody: HRP Goat Anti-rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blockipp byter 3% FSA

Blocking buffer: 3% BSA.



mmunohistochemistry of paraffin-embedded human preast carcinoma tissue, using Phospho-FOXO4-S197





Western blot analysis of various lysates, using Phospho-FOXO4-S197 antibody (STJ22147) at 1:40 dilution. Secondary antibody: HRP Goat Anti-Rabb (gG (H+L) (STJS000856) at 1:10000 dilutior; ysates/protiens: 25 Mu g per lane. Blocking buffer 3% nonfat dry milk in TBST. Detection: ECL Basic Kill Exposure fire. File.