

Anti-AMACR antibody (1-138) (STJ22599)
STJ22599

GENERAL INFORMATION


Product Type	Primary antibodies
Short Description	
Applications	WB/IHC-P/IF/ICC/ELISA
Host/Source	Rabbit
Reactivity	Human/Mouse/Rat

PRODUCT PROPERTIES


Clonality	Polyclonal
Clone ID	
Concentration	Lot specific
Conjugation	Unconjugated
Purification	Affinity purification
Dilution Range	WB:1:500-1:2000 IHC-P:1:50-1:200 IF/ICC:1:50-1:200 ELISA:Recommended starting concentration is 1 μ g/mL. Please optimize the concentration based on your specific assay requirements.
Formulation	PBS with 0.01% Thimerosal, 50% Glycerol, pH 7.3.
Isotype	IgG
Storage	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.
Instruction	

TARGET INFORMATION


Gene ID	23600
Gene Symbol	AMACR
Uniprot ID	AMACR_HUMAN
Immunogen	
Region	1-138
Specificity	Recombinant fusion protein containing a sequence corresponding to amino acids 1-138 of human AMACR (NP_055139.4).
Immunogen Sequence	MALQGISVVELSGLAPGPFC AMVLADFGARVVRVDRPGSR YDVSRLGRGKRSVLDLKQP RGAAVLRRLCKRSDVLLPEF RRGVMEKLQLQPEILQRENPLIYARLSGFGQSGSFCRLA GHDINYALSGVLSKIGR




Immunofluorescence analysis of mouse kidney using AMACR Rabbit polyclonal antibody (STJ22599) at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.



Immunofluorescence analysis of rat kidney using AMACR Rabbit polyclonal antibody (STJ22599) at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.



Immunohistochemistry analysis of paraffin-embedded human prostate cancer using AMACR Rabbit polyclonal antibody (STJ22599) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM Tris/EDTA buffer pH 9, 0 before commencing with immunohistochemistry staining protocol.



Western blot analysis of extracts of various cell lines, using AMACR antibody (STJ22599) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJ5000856) at 1:10000 dilution. Lysates/proteins: 25 μ g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 1s.

This product is suitable for in-vitro studies under the RESEARCH USE ONLY [RUO] licence. This product must not be used as for diagnostic or other medical purposes.
St John's Laboratory Ltd, Knowledge Dock Business Centre, University Way, London, E16 2RD | Tel: 0208 223 3081