

Anti-ANXA5 antibody (1-320) (STJ22620)
STJ22620

GENERAL INFORMATION


Product Type	Primary antibodies
Short Description	
Applications	WB/IF/ICC/ELISA
Host/Source	Rabbit
Reactivity	Human/Mouse

PRODUCT PROPERTIES


Clonality	Polyclonal
Clone ID	
Concentration	Lot specific
Conjugation	Unconjugated
Purification	Affinity purification
Dilution Range	WB:1:500-1:2000 IF/CC:1:50-1:200 ELISA:Recommended starting concentration is 1 Mu g/mL. Please optimize the concentration based on your specific assay requirements.
Formulation	PBS with 0.02% Sodium Azide, 50% Glycerol, pH 7.3.
Isotype	IgG
Storage Instruction	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

TARGET INFORMATION


Gene ID	308
Gene Symbol	ANXA5
Uniprot ID	ANXA5_HUMAN
Immunogen	
Immunogen Region	1-320
Specificity	Recombinant fusion protein containing a sequence corresponding to amino acids 1-320 of human Annexin A5/Annexin V (NP_001145.1).
Immunogen Sequence	MAQVLRGTVTDFPGFDERAD AETLRKAMKGLGTDEESILT LLTSRSNAQRQEISAFAKTL FGRDLLDLKSELTGKFEKL IVALMKPSRLYDAYELKHAL KGAGTNEKVLTEIASRTPE ELRAIKQVYEEYGGSSLEDD VVGDTSGYYQRMVLVLLQAN RDPDAGIDEAQVEQDAQALF QAGELKWTDEEKFITIFGT RSVSHLRKVFDKYMTISGFQ IETIDRETSGNLEQLLLA



Immunofluorescence analysis of HeLa cells using [KO Validated] Annexin A5/Annexin V Rabbit polyclonal antibody (STJ22620) at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.



Western blot analysis of extracts from normal (control) and Annexin A5/Annexin V knockout (KO) 293T cells, using Annexin A5/Annexin V antibody (STJ22620) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25 Mu g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 1s.



Western blot analysis of extracts of various cell lines, using Annexin A5/Annexin V antibody (STJ22620) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25 Mu g per lane. Blocking buffer: 3% nonfat dry milk in TBST.

This product is suitable for in-vitro studies under the RESEARCH USE ONLY [RUO] licence. This product must not be used as for diagnostic or other medical purposes.
St John's Laboratory Ltd, Knowledge Dock Business Centre, University Way, London, E16 2RD | Tel: 0208 223 3081