

Anti-CAPZA2 antibody (1-286) (STJ22888)

STJ22888

GENERAL INFORMATION


Product Type	Primary antibodies
Short Description	
Applications	WB/IHC-P/ELISA
Host/Source	Rabbit
Reactivity	Human/Mouse

PRODUCT PROPERTIES


Clonality	Polyclonal
Clone ID	
Concentration	Lot specific
Conjugation	Unconjugated
Purification	Affinity purification
Dilution	WB: 1:500-1:2000
Range	IHC-P: 1:50-1:200
	ELISA: Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.
Formulation	PBS with 0.09% Sodium Azide, 50% Glycerol, pH 7.3.
Isotype	IgG
Storage Instruction	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

TARGET INFORMATION


Gene ID	830
Gene Symbol	CAPZA2
Uniprot ID	CAZA2_HUMAN
Immunogen	
Immunogen Region	1-286
Specificity	Recombinant fusion protein containing a sequence corresponding to amino acids 1-286 of human CAPZA2 (NP_006127.1).
Immunogen Sequence	MADLEEQLSDEEKVRIAAKF IHHAPPGEFNEVFNDVRLLL NNDNLLREGAAHAFQYNLD QFTPVKIEGYEDQVLITEHG DLGNGKFLDPKNRICFKFDH LRKEATDPRPCEVENAVESW RTSVETALRAYVKEHYPNGV CTVYGKKIDGQQTIACIES HQFQAKNFWNGRWSEWKFT ITPSTTQVVGILKIQVHYE DGNVQLVSHKDIQDSLTVSN EVQTAKEFIKIVEAAENEY



Immunohistochemistry analysis of CAPZA2 in paraffin-embedded Mouse kidney using CAPZA2 Rabbit polyclonal antibody (STJ22888) at dilution of 1:50 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with immunohistochemistry staining protocol.



Immunohistochemistry analysis of CAPZA2 in paraffin-embedded Human colon using CAPZA2 Rabbit polyclonal antibody (STJ22888) at dilution of 1:50 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with immunohistochemistry staining protocol.



Western blot analysis of various lysates using CAPZA2 Rabbit polyclonal antibody (STJ22888) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJ5000856) at 1:10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST.

This product is suitable for in-vitro studies under the RESEARCH USE ONLY [RUO] licence. This product must not be used as for diagnostic or other medical purposes.
St John's Laboratory Ltd, Knowledge Dock Business Centre, University Way, London, E16 2RD | Tel: 0208 223 3081