

Anti-DCN antibody (31-359) (STJ23345)

STJ23345

GENERAL INFORMATION


Product Type	Primary antibodies
Short Description	
Applications	WB/IHC-P/IF/ICC/ELISA
Host/Source	Rabbit
Reactivity	Human/Mouse/Rat

PRODUCT PROPERTIES


Clonality	Polyclonal
Clone ID	
Concentration	Lot specific
Conjugation	Unconjugated
Purification	Affinity purification
Dilution	WB:1:500-1:1000
Range	IHC-P:1:50-1:200 IF/ICC:1:50-1:200 ELISA:Recommended starting concentration is 1 μ g/mL. Please optimize the concentration based on your specific assay requirements.
Formulation	PBS with 0.09% Sodium Azide, 50% Glycerol, pH 7.3.
Isotype	IgG
Storage	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.
Instruction	

TARGET INFORMATION


Gene ID	1634
Gene Symbol	DCN
Uniprot ID	PGS2_HUMAN
Immunogen	
Immunogen Region	31-359
Specificity	Recombinant fusion protein containing a sequence corresponding to amino acids 31-359 of Decorin (NP_001911.1).
Immunogen Sequence	DEASGIGPEVPDDRDFEPSL GPVCPFRQCCHLRVVQCSDL GLDKVPKDLPPDTLLDLQN NKITEIKDGDGFKNLKLNHAL ILVNNKISKVSPGAFTPLVK LERLYLSKNQKELPEKMPK TLQELRAHENEITKVRKVTF NGLNQMVIELGTNPLKSSG IENGAFQGMKKLSYIRIADT NITSIPQGLPPSLTELHLDG NKISRVDAASLKLNNLAKL GLSFNSISAVDNGSLANTP




Immunofluorescence analysis of MCF-7 cells using Decorin Rabbit polyclonal antibody (STJ23345). Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunohistochemistry analysis of Decorin in paraffin-embedded human colon using Decorin Rabbit polyclonal antibody (STJ23345) at dilution of 1:50 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with immunohistochemistry staining protocol.



Immunohistochemistry analysis of Decorin in paraffin-embedded human breast cancer using Decorin Rabbit polyclonal antibody (STJ23345) at dilution of 1:50 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with immunohistochemistry staining protocol.



Western blot analysis of lysates from Mouse heart, using Decorin Rabbit polyclonal antibody (STJ23345) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25 μ g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 180s.

This product is suitable for in-vitro studies under the RESEARCH USE ONLY [RUO] licence. This product must not be used as for diagnostic or other medical purposes.
St John's Laboratory Ltd, Knowledge Dock Business Centre, University Way, London, E16 2RD | Tel: 0208 223 3081