

Anti-NEDD1 antibody (411-660) (STJ24728)
STJ24728

GENERAL INFORMATION


Product Type	Primary antibodies
Short Description	
Applications	WB/IF/ICC/ELISA
Host/Source	Rabbit
Reactivity	Human/Mouse/Rat

PRODUCT PROPERTIES


Clonality	Polyclonal
Clone ID	
Concentration	Lot specific
Conjugation	Unconjugated
Purification	Affinity purification
Dilution	WB:1:500-1:2000
Range	IF/CC:1:50-1:200
	ELISA:Recommended starting concentration is 1 Mu g/mL. Please optimize the concentration based on your specific assay requirements.
Formulation	PBS with 0.02% Sodium Azide, 50% Glycerol, pH 7.3.
Isotype	IgG
Storage Instruction	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

TARGET INFORMATION

Gene ID	121441
Gene Symbol	NEDD1
Uniprot ID	NEDD1_HUMAN
Immunogen	
Immunogen Region	411-660
Specificity	Recombinant fusion protein containing a sequence corresponding to amino acids 411-660 of human NEDD1 (NP_690869.1).
Immunogen Sequence	SPIRDDAVVNGSGDESIGKG DGFDFLPQLNSVFPPRKNPV TSSTSVLHSSPLNVFMGSPG KEENENRDLTAESKKIYMGK QESKDSFKQLAKLVTSGAES GNLNTSPSSNQTRNSEKFEK PENEIEAQLICEPPINGSST PNPKIASSVTAGVASSLSEK IADSIGNNRQNAPLTSIQIR FIQNMIQETLDDFREACHRD IVNLQVEMIKQFHMQLNEMH SLLERYSVNEGLVAEIERL



Immunofluorescence analysis of L929 cells using NEDD1 antibody (STJ24728) at dilution of 1:100. Blue: DAPI for nuclear staining.



Western blot analysis of extracts of various cell lines, using NEDD1 antibody (STJ24728) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJ3000896) at 1:10000 dilution. Lysates/proteins: 25 Mu g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 30s.

This product is suitable for in-vitro studies under the RESEARCH USE ONLY [RUO] licence. This product must not be used as for diagnostic or other medical purposes.
St John's Laboratory Ltd, Knowledge Dock Business Centre, University Way, London, E16 2RD | Tel: 0208 223 3081