

Anti-NLK antibody (278-527) (STJ24772)
STJ24772

GENERAL INFORMATION


Product Type	Primary antibodies
Short Description	
Applications	WB/IHC-P/ELISA
Host/Source	Rabbit
Reactivity	Human/Mouse/Rat

PRODUCT PROPERTIES


Clonality	Polyclonal
Clone ID	
Concentration	Lot specific
Conjugation	Unconjugated
Purification	Affinity purification
Dilution Range	WB:1:500-1:2000 IHC-P:1:50-1:200 ELISA:Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.
Formulation	PBS with 0.02% Sodium Azide, 50% Glycerol, pH 7.3.
Isotype	IgG
Storage Instruction	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

TARGET INFORMATION


Gene ID	51701
Gene Symbol	NLK
Uniprot ID	NLK_HUMAN
Immunogen	
Immunogen Region	278-527
Specificity	Recombinant fusion protein containing a sequence corresponding to amino acids 278-527 of human NLK (NP_057315.3).
Immunogen Sequence	LKICDFGLARVEELDESRHM TQEVVTQYYRAPEILMGSRH YSNAIDIWSVGCIFAELLGR RILFQAQSPIQQDLITDLL GTPSLEAMRTACEGAHAHIL RGPHEKQPSLPVLYTLSSQAT HEAVHLLCRMLVFDPSKRIS AKDALAHPLYLDEGRRLRYHTC MCKCCFSTSTGRVYTSDFEP VTNPKFDDTFEKNLSSVRQV KEIIHQFILEQQKGNRVPLC INPQSAAFKSFISSTVAQP




Immunohistochemistry analysis of paraffin-embedded rat liver using NLK Rabbit polyclonal antibody (STJ24772) at dilution of 1:20 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with immunohistochemistry staining protocol.



Immunohistochemistry analysis of paraffin-embedded mouse kidney using NLK Rabbit polyclonal antibody (STJ24772) at dilution of 1:20 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with immunohistochemistry staining protocol.



Immunohistochemistry analysis of paraffin-embedded human liver cancer using NLK Rabbit polyclonal antibody (STJ24772) at dilution of 1:20 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with immunohistochemistry staining protocol.



This product is suitable for in-vitro studies under the RESEARCH USE ONLY [RUO] licence. This product must not be used as for diagnostic or other medical purposes.
St John's Laboratory Ltd, Knowledge Dock Business Centre, University Way, London, E16 2RD | Tel: 0208 223 3081