

## Anti-TSPAN33 antibody (C-Term) (STJ27195)

STJ27195

### GENERAL INFORMATION


|                          |                    |
|--------------------------|--------------------|
| <b>Product Type</b>      | Primary antibodies |
| <b>Short Description</b> |                    |
| <b>Applications</b>      | WB/ELISA           |
| <b>Host/Source</b>       | Rabbit             |
| <b>Reactivity</b>        | Human/Mouse/Rat    |

### PRODUCT PROPERTIES

|                            |                                                                                                                                                      |
|----------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Clonality</b>           | Polyclonal                                                                                                                                           |
| <b>Clone ID</b>            |                                                                                                                                                      |
| <b>Concentration</b>       | Lot specific                                                                                                                                         |
| <b>Conjugation</b>         | Unconjugated                                                                                                                                         |
| <b>Purification</b>        | Affinity purification                                                                                                                                |
| <b>Dilution Range</b>      | WB:1:200-1:1000<br>ELISA:Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements. |
| <b>Formulation</b>         | PBS with 0.01% Thimerosal, 50% Glycerol, pH 7.3.                                                                                                     |
| <b>Isotype</b>             | IgG                                                                                                                                                  |
| <b>Storage Instruction</b> | Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.                                                       |

### TARGET INFORMATION

|                           |                                                                                                                          |
|---------------------------|--------------------------------------------------------------------------------------------------------------------------|
| <b>Gene ID</b>            | 340348                                                                                                                   |
| <b>Gene Symbol</b>        | TSPAN33                                                                                                                  |
| <b>Uniprot ID</b>         | TSN33_HUMAN                                                                                                              |
| <b>Immunogen</b>          |                                                                                                                          |
| <b>Immunogen Region</b>   | C-Term                                                                                                                   |
| <b>Specificity</b>        | A synthetic peptide corresponding to a sequence within amino acids 200 to the C-terminus of human TSPAN33 (NP_848657.1). |
| <b>Immunogen Sequence</b> | NTMCGQGGMQAFDYLEASKVI YTNCGIDKLVNWIHSNLFLL GGVALGLAIPQLVGILLSQI LVNQIKDQIKLQLYNQQHRA DPWY                                |



Western blot analysis of various lysates using TSPAN33 Rabbit polyclonal antibody (STJ27195) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 90s.

This product is suitable for in-vitro studies under the RESEARCH USE ONLY [RUO] licence. This product must not be used as for diagnostic or other medical purposes.  
St John's Laboratory Ltd, Knowledge Dock Business Centre, University Way, London, E16 2RD | Tel: 0208 223 3081