

Anti-ARG2 antibody (270-354) (STJ28438)

STJ28438

GENERAL INFORMATION


Product Type	Primary antibodies
Short Description	
Applications	WB/ELISA
Host/Source	Rabbit
Reactivity	Human/Mouse/Rat

PRODUCT PROPERTIES


Clonality	Polyclonal
Clone ID	
Concentration	Lot specific
Conjugation	Unconjugated
Purification	Affinity purification
Dilution Range	WB:1:500-1:1000 ELISA:Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.
Formulation	PBS with 0.05% Proclin300, 50% Glycerol, pH 7.3.
Isotype	IgG
Storage Instruction	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

TARGET INFORMATION

Gene ID	384
Gene Symbol	ARG2
Uniprot ID	ARGI2_HUMAN
Immunogen	
Immunogen Region	270-354
Specificity	Recombinant fusion protein containing a sequence corresponding to amino acids 270-354 of human Arginase 2 (ARG2) (NP_001163.1).
Immunogen Sequence	GLTYREGMYIAEEIHNTGLL SALDLVEVNPQLATSEEEAK TTANLAVDVIASSFGQTREG GHIVYDQLPTPSPDESENQ ARVRI



Western blot analysis of extracts of Rat kidney, using Arginase 2 (ARG2) antibody (STJ28438) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 180s.



Western blot analysis of extracts of various cell lines, using Arginase 2 (ARG2) antibody (STJ28438) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 1s.

This product is suitable for in-vitro studies under the RESEARCH USE ONLY [RUO] licence. This product must not be used as for diagnostic or other medical purposes.
St John's Laboratory Ltd, Knowledge Dock Business Centre, University Way, London, E16 2RD | Tel: 0208 223 3081