

Anti-RPS14 antibody (1-151) (STJ28810)
STJ28810

GENERAL INFORMATION


Product Type	Primary antibodies
Short Description	
Applications	WB/IHC-P/IF/ICC/ELISA
Host/Source	Rabbit
Reactivity	Human/Mouse/Rat

PRODUCT PROPERTIES


Clonality	Polyclonal
Clone ID	
Concentration	Lot specific
Conjugation	Unconjugated
Purification	Affinity purification
Dilution	WB:1:500-1:2000
Range	IHC-P:1:50-1:200 IF/ICC:1:50-1:200 ELISA:Recommended starting concentration is 1 Mu g/mL. Please optimize the concentration based on your specific assay requirements.
Formulation	PBS with 0.02% Sodium Azide, 50% Glycerol, pH 7.3.
Isotype	IgG
Storage	Store at-20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.
Instruction	

TARGET INFORMATION


Gene ID	6208
Gene Symbol	RPS14
Uniprot ID	RS14_HUMAN
Immunogen	1-151
Region	
Specificity	Recombinant fusion protein containing a sequence corresponding to amino acids 1-151 of human RPS14 (NP_005608.1).
Immunogen	MAPRKKGKEKKEEQVISLGPQ VAEGENVFGVCHIFASFNDT FVHVTDLSGKETICRVTGGM KVKADRDESSPYAAMLAQD
Sequence	VAQRCKELGITALHIKLRAT GGNRTKTPGPGAQSALRALA RSGMGKIGRIEDVTPIPSDST RRKGRRGRRL




Immunofluorescence analysis of MCF7 cells using RPS14 antibody (STJ28810). Blue: DAPI for nuclear staining.



Immunohistochemistry analysis of paraffin-embedded mouse spinal cord using RPS14 Rabbit polyclonal antibody (STJ28810) at dilution of 1:200 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with immunohistochemistry staining protocol.



Immunohistochemistry analysis of paraffin-embedded rat pancreas using RPS14 Rabbit polyclonal antibody (STJ28810) at dilution of 1:200 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with immunohistochemistry staining protocol.



Western blot analysis of extracts of various cell lines, using RPS14 antibody (STJ28810) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJ5000856) at 1:10000 dilution. Lysates/proteins: 25 Mu g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 30s.

This product is suitable for in-vitro studies under the RESEARCH USE ONLY [RUO] licence. This product must not be used as for diagnostic or other medical purposes.
St John's Laboratory Ltd, Knowledge Dock Business Centre, University Way, London, E16 2RD | Tel: 0208 223 3081