

Anti-TRAF3IP2 antibody (1-300) (STJ28859)
STJ28859

GENERAL INFORMATION


Product Type	Primary antibodies
Short Description	
Applications	WB/IF/ICC/ELISA
Host/Source	Rabbit
Reactivity	Human/Mouse/Rat

PRODUCT PROPERTIES


Clonality	Polyclonal
Clone ID	
Concentration	Lot specific
Conjugation	Unconjugated
Purification	Affinity purification
Dilution Range	WB:1:500-1:1000 IF/ICC:1:50-1:200 ELISA:Recommended starting concentration is 1 Mu g/mL. Please optimize the concentration based on your specific assay requirements.
Formulation	PBS with 0.02% Sodium Azide, 50% Glycerol, pH 7.3.
Isotype	IgG
Storage Instruction	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

TARGET INFORMATION


Gene ID	10758
Gene Symbol	TRAF3IP2
Uniprot ID	CIKS_HUMAN
Immunogen	
Immunogen Region	1-300
Specificity	Recombinant fusion protein containing a sequence corresponding to amino acids 1-300 of human TRAF3IP2 (NP_671733.2).
Immunogen Sequence	MPPQLQETRMNRSIPVEVDE SEPYPSQLLKPIEYSPEEE SEPPAPNIRNMAPNSLSAPT MLHNSSGDFSQAHSTLKLAN HQRPVSRQVTCLRTQVLEDS EDSFCRRHPGLGKAFPSGCS AVSEPASESVVGALPAEHQF SFMEKRNQWLVSQLSAASPD TGHDSKSDQSLPNASADSL GGSQEMVQRPPQHRNRAGLD LPTIDTGYDSQPQDVLGIRQ LERPLPLTSVCYPQDLPRP



Immunofluorescence analysis of Hep G2 cells using TRAF3IP2 Rabbit polyclonal antibody (STJ28859) at a dilution of 1:200 (40x lens). Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) at 1:500 dilution. Blue: DAPI for nuclear staining.



Western blot analysis of lysates from Rat lung, using TRAF3IP2 Rabbit polyclonal antibody (STJ28859) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25 Mu g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 90s.



Western blot analysis of various lysates using TRAF3IP2 Rabbit polyclonal antibody (STJ28859) at 1:500 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25 Mu g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 180s.

This product is suitable for in-vitro studies under the RESEARCH USE ONLY [RUO] licence. This product must not be used as for diagnostic or other medical purposes.
St John's Laboratory Ltd, Knowledge Dock Business Centre, University Way, London, E16 2RD | Tel: 0208 223 3081