

Anti-PTGES2 antibody (88-377) (STJ29217)

STJ29217

GENERAL INFORMATION


Product Type	Primary antibodies
Short Description	
Applications	WB/IHC-P/IF/ICC/ELISA
Host/Source	Rabbit
Reactivity	Human/Mouse/Rat

PRODUCT PROPERTIES


Clonality	Polyclonal
Clone ID	
Concentration	Lot specific
Conjugation	Unconjugated
Purification	Affinity purification
Dilution	WB:1:500-1:2000
Range	IHC-P:1:50-1:200 IF/ICC:1:50-1:100 ELISA:Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.
Formulation	PBS with 0.02% Sodium Azide, 50% Glycerol, pH 7.3.
Isotype	IgG
Storage	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.
Instruction	

TARGET INFORMATION


Gene ID	80142
Gene Symbol	PTGES2
Uniprot ID	PGES2_HUMAN
Immunogen	
Immunogen Region	88-377
Specificity	Recombinant fusion protein containing a sequence corresponding to amino acids 88-377 of human PTGES2 (NP_079348.1).
Immunogen Sequence	ERSAAQLSLSSRLQLTYQY KTCPFCSKVRAFLDFHALPY QVVEVNPVRRAEIKFSSYRK VPILVAQEGESSQQLNDSSV IISALKTYLVSGQPLEEIT YYPAMKAVNEQGKEVTEFGN KYWLMMLNEKAQQVYGGKEA RTEEMKWRQWADDWLVLHLIS PNVYRTPTALASFDYIVRE GKFGAVEGAVAKYMGAAAMY LISRLKSRHRLQDNVREDL YEAADKWWAAVVGKDRPFMGM




Immunofluorescence analysis of A-549 cells using PTGES2 antibody (STJ29217). Blue: DAPI for nuclear staining.



Immunohistochemistry analysis of paraffin-embedded mouse kidney using PTGES2 antibody (STJ29217) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with immunohistochemistry staining protocol.



Immunohistochemistry analysis of paraffin-embedded human kidney cancer using PTGES2 antibody (STJ29217) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with immunohistochemistry staining protocol.



Immunohistochemistry analysis of paraffin-embedded rat kidney using PTGES2 antibody (STJ29217) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with immunohistochemistry staining protocol.

This product is suitable for in-vitro studies under the RESEARCH USE ONLY [RUO] licence. This product must not be used as for diagnostic or other medical purposes.
St John's Laboratory Ltd, Knowledge Dock Business Centre, University Way, London, E16 2RD | Tel: 0208 223 3081