Anti-Phospho-BCR-Y177 antibody (STJ29372)

STJ29372

GENERAL INFORMATION

Product Type Primary antibodies
Short Description
Applications WB/ELISA
Host/Source Rabbit
Reactivity Human/Mouse

PRODUCT PROPERTIES

Clonality Polyclonal
Clone ID
Concentration Lot specific
Conjugation Unconjugated
Purification Affinity purification
Dilution Range WB:1:500-1:2000
ELISA:Recommended starting concentration is 1 Mu g/mL. Please optimize the concentration based on your specific assay requirements.
Formulation PBS with 0.02% Sodium Azide, 50% Glycerol, pH 7.3.
Isotype IgG
Storage Instruction Store at-20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

TARGET INFORMATION

Gene ID 613
Gene Symbol BCR
Uniprot ID BCR_HUMAN
Immunogen PFYVN
Immunogen Region
Specificity A synthetic phosphorylated peptide around Y177 of human BCR (NP_004318.3).
Immunogen Sequence PFYVN



Western blot analysis of extracts of K562 cells, using Phospho-BCR-Y177 antibody (STJ29372) at 1:1000 dilution. K562 cell lysate were treated by CIP (20ul CIP for each 400ul cell lysate) at 37 Â°C for 1 hour. Secondary antibody: HRP Goat Anti-rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% BSA. Detection: ECL Basic Kit. Exposure time: 90s.



Western blot analysis of extracts of K562 cells, using Phospho-BCR-Y177 antibody (STJ29372) at 1:1000 dilution. K562 cell lysate were treated by CIP (20ul CIP for each 400ul cell lysate) at 37 Â°C for 1 hour. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25 Mu g per lane. Blocking buffer: 3% BSA. Detection: ECL Basic Kit. Exposure time: 90s.

St John's Laboratory Ltd, Knowledge Dock Business Centre, University Way, London, E16 2RD | Tel: 0208 223 3081