

Anti-MID1 antibody (478-667) (STJ29430)
STJ29430

GENERAL INFORMATION


Product Type	Primary antibodies
Short Description	
Applications	WB/IF/ICC/ELISA
Host/Source	Rabbit
Reactivity	Human

PRODUCT PROPERTIES


Clonality	Polyclonal
Clone ID	
Concentration	Lot specific
Conjugation	Unconjugated
Purification	Affinity purification
Dilution	WB:1:500-1:2000
Range	IF/CC:1:50-1:100 ELISA:Recommended starting concentration is 1 Mu g/mL. Please optimize the concentration based on your specific assay requirements.
Formulation	PBS with 0.02% Sodium Azide, 50% Glycerol, pH 7.3.
Isotype	IgG
Storage Instruction	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

TARGET INFORMATION

Gene ID	4281
Gene Symbol	MID1
Uniprot ID	TR18_HUMAN
Immunogen	
Immunogen Region	478-667
Specificity	Recombinant fusion protein containing a sequence corresponding to amino acids 478-667 of human MID1 (NP_000372.1).
Immunogen Sequence	KLKTN SQPFKLPKSAHRKL KVSHDNLTVERDESSSKKSH TPERFTSQGSYG VAGNVFID SGRHYWEVVISGSTWY AIGL AYKSAPKHEWIGKNSASWAL CRCNNNWVVRHNSKEIPIEP APHLRRV GILLDYDNGSIAF YDALNSIHLYTFDVAFAQP V CPTFTVWNKCLTIITGLPIP DHLDCTEQLP



Immunofluorescence analysis of MCF-7 cells using MID1 Rabbit polyclonal antibody (STJ29430). Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) at 1:500 dilution. Blue: DAPI for nuclear staining.



Western blot analysis of lysates from HeLa cells, using MID1 Rabbit polyclonal antibody (STJ29430) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJ5000856) at 1:10000 dilution. Lysates/proteins: 25 Mu g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 60s.

This product is suitable for in-vitro studies under the RESEARCH USE ONLY [RUO] licence. This product must not be used as for diagnostic or other medical purposes.
St John's Laboratory Ltd, Knowledge Dock Business Centre, University Way, London, E16 2RD | Tel: 0208 223 3081