

Anti-DFFA antibody (1-331) (STJ29835)

STJ29835

GENERAL INFORMATION


Product Type	Primary antibodies
Short Description	
Applications	WB/IF/ICG/ELISA
Host/Source	Rabbit
Reactivity	Human/Mouse/Rat

PRODUCT PROPERTIES


Clonality	Polyclonal
Clone ID	
Concentration	Lot specific
Conjugation	Unconjugated
Purification	Affinity purification
Dilution	WB:1:500-1:2000
Range	IF/ICG:1:50-1:100 ELISA:Recommended starting concentration is 1 Mu g/mL. Please optimize the concentration based on your specific assay requirements.
Formulation	PBS with 0.02% Sodium Azide, 50% Glycerol, pH 7.3.
Isotype	IgG
Storage Instruction	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

TARGET INFORMATION


Gene ID	1676
Gene Symbol	DFFA
Uniprot ID	DFFA_HUMAN
Immunogen	
Immunogen Region	1-331
Specificity	Recombinant fusion protein containing a sequence corresponding to amino acids 1-331 of human DFFA (NP_004392.1).
Immunogen Sequence	MEVTGDAGVPESGEIRTLKP CLLRRNYSREQHGVAASCLE DLRKACDILAIKSLTPVT LVLAEDGTIVDDDDYFLCLP SNTKFFVALASNEKWAYNNNSD GGTAWISQESFDVDETDGSA GLKWKNVARQLKEDLSSIIL LSEEDLQMLVDAPCSDLAQE LRQSCATVRLQHTLQQVLD QREEVRQSKQLLQYLQALE KEGSLLSKQESKAAFGEV DAVDTGISRETSSDVALAS



Immunofluorescence analysis of HeLa cells using [KO Validated] DFFA Rabbit polyclonal antibody (STJ29835). Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) at 1:500 dilution. Blue: DAPI for nuclear staining.



Western blot analysis of lysates from wild type (WT) and DFFA knockout (KO) HeLa cells, using [KO Validated] DFFA Rabbit polyclonal antibody (STJ29835) at 1:3000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJ5000856) at 1:10000 dilution. Lysates/proteins: 25 Mu g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 5s.



Western blot analysis of various lysates using [KO Validated] DFFA Rabbit polyclonal antibody (STJ29835) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJ5000856) at 1:10000 dilution. Lysates/proteins: 25 Mu g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 15s.

This product is suitable for in-vitro studies under the RESEARCH USE ONLY [RUO] licence. This product must not be used as for diagnostic or other medical purposes.

St John's Laboratory Ltd, Knowledge Dock Business Centre, University Way, London, E16 2RD | Tel: 0208 223 3081