

## Anti-KHDRBS1 antibody (343-443) (STJ29932)

STJ29932

### GENERAL INFORMATION


<b>Product Type</b>	Primary antibodies
<b>Short Description</b>	
<b>Applications</b>	WB/IHC-P/IF/ICC/ELISA
<b>Host/Source</b>	Rabbit
<b>Reactivity</b>	Human/Mouse/Rat

### PRODUCT PROPERTIES


<b>Clonality</b>	Polyclonal
<b>Clone ID</b>	
<b>Concentration</b>	Lot specific
<b>Conjugation</b>	Unconjugated
<b>Purification</b>	Affinity purification
<b>Dilution Range</b>	WB:1:500-1:2000 IHC-P:1:100-1:200 IF/ICC:1:50-1:200 ELISA:Recommended starting concentration is 1 Mu g/mL. Please optimize the concentration based on your specific assay requirements.
<b>Formulation</b>	PBS with 0.02% Sodium Azide, 50% Glycerol, pH 7.3.
<b>Isotype</b>	IgG
<b>Storage Instruction</b>	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

### TARGET INFORMATION


<b>Gene ID</b>	10657
<b>Gene Symbol</b>	KHDRBS1
<b>Uniprot ID</b>	KHDR1_HUMAN
<b>Immunogen</b>	
<b>Immunogen Region</b>	343-443
<b>Specificity</b>	A synthetic peptide corresponding to a sequence within amino acids 343-443 of human KHDRBS1/Sam68 (NP_006550.1).
<b>Immunogen Sequence</b>	PAPRARTAGIQRILPPPPA PETYEYGYDDTYAEQSYEG YEGYYSQSQGDSEYYDYGHG EVQDSYEAYGQDDWNGTRPS LKAPPARPVKGAYREHPYGR Y




Confocal immunofluorescence analysis of U2OS cells using KHDRBS1/Sam68 Polyclonal Antibody (STJ29932) at dilution of 1:400. Blue: DAPI for nuclear staining.



Confocal immunofluorescence analysis of HeLa cells using KHDRBS1/Sam68 Polyclonal Antibody (STJ29932) at dilution of 1:400. Blue: DAPI for nuclear staining.



Immunohistochemistry analysis of paraffin-embedded mouse liver using KHDRBS1/Sam68 antibody (STJ29932) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with immunohistochemistry staining protocol.



Immunohistochemistry analysis of paraffin-embedded human stomach using KHDRBS1/Sam68 antibody (STJ29932) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with immunohistochemistry staining protocol.

This product is suitable for in-vitro studies under the RESEARCH USE ONLY [RUO] licence. This product must not be used as for diagnostic or other medical purposes.  
St John's Laboratory Ltd, Knowledge Dock Business Centre, University Way, London, E16 2RD | Tel: 0208 223 3081