

Anti-TANK2 antibody (Internal) (STJ70000)

STJ70000

GENERAL INFORMATION


Product Type	Primary antibodies
Short Description	Goat polyclonal antibody anti-TANK2 (Internal) is suitable for use in Western Blot research applications.
Applications	WB-Trf
Host/Source	Goat
Reactivity	Human/Mouse/Rat/Dog

PRODUCT PROPERTIES

Clonality	Polyclonal
Clone ID	
Concentration	0.5 mg/mL
Conjugation	Unconjugated
Purification	Purified from goat serum by ammonium sulphate precipitation followed by affinity chromatography using Protein G.
Dilution Range	WB: In transfected HEK293 transiently expressing TANK2 a band of approx. 127kDa is observed. This band is not observed in the non-transfected HEK293. The calculated molecular size is 127 kDa according to NP_079511.1. Recommended concentration, 0.5-1.
Formulation	0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. NA
Isotype	IgG
Storage Instruction	Store at -20°C on receipt and minimise freeze-thaw cycles.

TARGET INFORMATION

Gene ID	80351
Gene Symbol	TNKS2
Uniprot ID	TNKS2_HUMAN
Immunogen	
Immunogen Region	Internal
Specificity	Fusion protein containing 73 aas from region aa800-900 (according to NP_079511.1). No crossreactivity expected with TANK1.
Immunogen Sequence	RECOMBINANT2



This product is suitable for in-vitro studies under the RESEARCH USE ONLY [RUO] licence. This product must not be used as for diagnostic or other medical purposes.
St John's Laboratory Ltd, Knowledge Dock Business Centre, University Way, London, E16 2RD | Tel: 0208 223 3081