

## Anti-FGR antibody (C-Term) (STJ70074)

ST.170074

## **GENERAL INFORMATION**

Product Type Primary antibodies

Short Goat polyclonal antibody anti-FGR (C-Term) is suitable for use in ELISA and Western Blot research applications.

Description

Applications Pep-ELISA/WB

Host/Source Goat

Reactivity Human/Mouse/Rat/Dog

## **PRODUCT PROPERTIES**

Clonality Polyclonal
Clone ID

Concentration 0.5 mg/mL
Conjugation Unconjugated

Purification Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing

peptide.

**Dilution Range** Peptide ELISA: antibody detection limit dilution 1:128000.

WB: Approx 60kDa band observed in Human peripheral blood monocytes lysate (calculated MW of 59.5kDa according to

NP\_001036194.1). Recommended concentration: 0.3-1µg/ml.

 $\textbf{Formulation} \quad 0.5 \text{ mg/ml in Tris saline, } 0.02\% \text{ sodium azide, pH7.3 with } 0.5\% \text{ bovine serum albumin. NA}$ 

Isotype IgG

Storage Store at-20°C on receipt and minimise freeze-thaw cycles.

Instruction

## **TARGET INFORMATION**

 Gene ID
 2268

 Gene Symbol
 FGR

 Uniprot ID
 FGR\_HUMAN

 Immunogen
 C-Term

Region

Specificity This antibody is expected to recognise all three reported isoforms (NP\_001036194.1; NP\_001036212.1; NP\_005239.1).

Immunogen TSAEPQYQPGDQT

Sequence

250kDa 150kDa 100kDa 75kDa 50kDa 37kDa 25kDa 20kDa

STJ70074 (0. 3ŵg/ml) staining of Human periphera blood lymphocytes lysate (35ŵg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.