

Anti-DAP3 antibody (C-Term) (STJ70171)

GENERAL INFORMATION

Product Type Primary antibodies

Short Goat polyclonal antibody anti-DAP3 (C-Term) is suitable for use in ELISA, Western Blot, Immunofluorescence and

Description Immunohistochemistry research applications.

Applications Pep-ELISA/WB/IF/IHC

Host/Source Goat Reactivity Human

PRODUCT PROPERTIES

Clonality Polyclonal

Clone ID Concentration 0.5 mg/mL Conjugation Unconjugated

Purification Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing

Dilution Peptide ELISA: antibody detection limit dilution 1:64000.

Range WB: Approx 40kDa band observed in lysates of cell lines HeLa and HepG2 and in Human Kidney lysates (calculated MW of 41kDa

according to NP_001186779.1 and NP_001186780). Recommended concen

 $\textbf{Formulation} \quad 0.5 \text{ mg/ml in Tris saline, } 0.02\% \text{ sodium azide, pH7.3 with } 0.5\% \text{ bovine serum albumin. NA}$

Isotype IgG

Storage Store at-20°C on receipt and minimise freeze-thaw cycles.

Instruction

TARGET INFORMATION

Gene ID 7818 Gene Symbol DAP3

Immunogen

Uniprot ID RT29_HUMAN

Immunogen C-Term


Region Specificity


This antibody is expected to recognise isoform 1 (NP_387506.1), isoform 2 (NP_001186779.1) and isoform 3 (NP_001186780.1).


Reported variants represent identical protein (NP_387506.1; NP_004623.1; NP_001186778.1).


Immunogen NPSLLERHCAYL

Sequence









STJ70171 (1ŵg/ml) staining of HeLa lysate (control in left lane and after si-RNA-mediated DAP3 knock-down expresson in right lane) S5Aµg protein in RIPA buffer). Level of knock-down relative to Actin expression level was determined by RT-PCR. Primary inculpation was 1