Anti-Dynactin antibody (C-Term) (STJ70251)
STJ70251

GENERAL INFORMATION

Product Type	Primary antibodies
Short Description	Goat polyclonal antibody anti-Dynactin (C-Term) is suitable for use in ELISA, Western Blot, Immunofluorescence, Immunohistochemistry and Flow Cytometry research applications.
Applications	Pep-ELISA/WB/IF/IHC/FC
Host/Source	Goat
Reactivity	Human/Mouse

PRODUCT PROPERTIES

Clonality	Polyclonal
Clone ID	
Concentration	0.5 mg/mL
Conjugation	Unconjugated
Purification	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
Dilution Range	Peptide ELISA: antibody detection limit dilution 1:16000. WB: Approx. 150kDa band observed in HeLa and MCF7 cell lysates, and in preliminary testing of HEK293 and Mouse Brain lysate (calculated MW of 142kDa according to NP_004073.2). Recommended
Formulation	0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. NA
Isotype	IgG
Storage Instruction	Store at-20°C on receipt and minimise freeze-thaw cycles.

TARGET INFORMATION

Gene ID	1639
Gene Symbol	DCTN1
Uniprot ID	DCTN1_HUMAN
Immunogen	
Immunogen Region	C-Term
Specificity	This antibody is expected to recognise all reported isoforms.
Immunogen Sequence	QEQLHQLHSRLIS

STJ70251 (1µg/ml) staining of Human Cerebellum lysate (35µg protein in RIPA buffer). Detected by chemiluminescence.

STJ70251 Immunofluorescence analysis of paraformaldehyde fixed HeLa cells, permeabilized with 0. 15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml) , showing cytoplasmic staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).

STJ70251 Immunofluorescence analysis of paraformaldehyde fixed U2OS cells, permeabilized with 0. 15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml) , showing cytoplasmic and nuclear staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).

STJ70251 Flow cytometric analysis of paraformaldehyde fixed HeLa cells (blue line) , permeabilized with 0. 5% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (1ug/ml). IgG control: Unimmunized goat IgG (black line) followed by Alexa Fluor 488 secondary antibody.